

Regulatory Roles of MicroRNAs in Female and Male Human Reproduction: A Narrative Review

Mohadeseh Khoshandam^{1,2}, PhD;
Leila Naserpour^{1,3}, PhD; Mohammad
Taghi Hedayati Goudarzi⁴, MD; Hossein
Soltaninejad⁵, PhD; Hadi Zare-Zardini⁶,
PhD; Ashraf Alsadat Hoseiny¹, MSc; Neda
Golpar Raboki¹, MSc; Elahe Sadat Mousavi¹,
MSc; Mohammad Ali Salami⁷, PhD

¹Department of Reproductive Biology,
Academic Center for Education, Culture,
and Research (ACECR), Qom, Iran;

²National Institute of Genetic Engineering
and Biotechnology (NIGEB), Tehran, Iran;

³Department of Tissue Engineering, Qom
University of Medical Sciences, Qom, Iran;

⁴Department of Cardiology, School of
Medicine, Babol University of Medical
Sciences, Babol, Iran;

⁵Department of Stem Cells Technology
and Tissue Regeneration, Faculty of
Interdisciplinary Science and Technologies,
Tarbiat Modares University, Tehran, Iran;

⁶Department of Biomedical Engineering,
Meybod University, Meybod, Iran;

⁷Department of Biomaterials,
Nanotechnology and Tissue Engineering,
School of Advanced Technologies in
Medicine, Isfahan University of Medical
Sciences, Isfahan, Iran

Correspondence:

Hossein Soltaninejad, PhD;
Tarbiat Modares University, Jalal Al Ahmad
Hwy., Nasr Bridge, Postal code: 14117-13116,
Tehran, Iran

Email: hosoltaninejad@gmail.com
hossein.soltani@modares.ac.ir

Received: 12 November 2024

Revised: 16 October 2024

Accepted: 23 December 2024

What's Known

- The role of microRNAs (miRNAs) in human reproduction is an exciting area of research, and these regulatory molecules likely play an essential role in reproductive function. However, the current understanding of miRNAs' regulatory role in male and female reproduction remains unclear, with a considerable amount of conflicting and inconsistent data.

What's New

- An extensive review of studies revealed a consistent relationship between aberrant miRNA expression and infertility. Therefore, measuring the expression of specific miRNAs identified in this review could serve as valuable biomarkers for male and female infertility and could lead to the development of novel and specialized treatments.

Abstract

The role of microRNAs (miRNAs) in human reproduction represents an area of research, as these regulatory molecules appear to play essential roles in reproductive function. However, the current understanding of miRNAs' regulatory mechanisms in male and female reproduction remains incomplete, with considerable contradictory evidence in the literature. This targeted review aimed to analyze high-quality studies published to date on miRNA expression patterns in female and male reproduction to elucidate their biological roles and associations with infertility, thereby updating knowledge in this field. A comprehensive review of the literature was conducted using different electronic databases and search engines, including PubMed Central, MEDLINE, and Google Scholar from the earliest available records up to 2023. This search identified 18,100 articles related to miRNA expression in infertility, polycystic ovary syndrome (PCOS), endometriosis, and biomarkers, of which 72 met our criteria for further analysis. The findings of the present study revealed that specific, stable miRNA populations exist in different tissues and cells, potentially influencing spermatogenesis and oogenesis. The extensive review of studies suggested a consistent relationship between aberrant miRNA expression patterns and infertility. Consequently, the miRNAs identified in this review might serve as valuable biomarkers for both male and female infertility and could lead to the development of novel and specialized treatments.

This manuscript has been released as a preprint at Authorea: (<https://doi.org/10.22541/au.169609839.97787658/v1>).

Please cite this article as: Khoshandam M, Naserpour L, Hedayati Goudarzi MT, Soltaninejad H, Zare-Zardini H, Hoseiny AA, Golpar Raboki N, Mousavi ES, Salami MA. Regulatory Roles of MicroRNAs in Female and Male Human Reproduction: A Narrative Review. Iran J Med Sci. 2025;50(9):588-596. doi: 10.30476/ijms.2025.104375.3797.

Keywords • Infertility • MicroRNAs • Spermatogenesis • Polycystic ovary syndrome • Endometriosis • Biomarkers

Introduction

MicroRNAs (miRNAs) are single-stranded RNA molecules approximately 22 nucleotides long.¹ These molecules play a significant and crucial role in gene expression through partial complementarity with the 3' untranslated region (3'UTR).² Several miRNAs have been identified in humans, and research has shown their importance in biological processes, including cell cycle regulation, differentiation, metabolism, proliferation, apoptosis, gametogenesis (both male and female), and embryo development.³

Over the past decade, studies have indicated the ubiquitous

presence of miRNAs across all human tissues, including the reproductive system.^{4, 5} The identification of miRNAs in various reproductive organs and tissues, such as the testis, ovary, uterus, epididymis, spermatozoa, oocytes, seminal plasma, and extracellular vesicles, coupled with their known regulatory functions, suggested that miRNA dysregulation might impair spermatogenesis, disrupt oocyte maturation, and contribute to various types of infertility.⁶

This study aimed to systematically review high-quality published studies to investigate the association between miRNAs and reproductive functions in both sexes, with particular focus on their role in infertility. While the available studies demonstrated considerable heterogeneity, a comprehensive synthesis of this evidence could help consolidate findings, resolve existing inconsistencies, and provide guidance for future research.

The growing urgency for rapid infertility diagnostics and treatments highlighted the potential of innovative approaches, such as liquid biopsy, which offers faster and less invasive alternatives to conventional approaches. Given their established role in gene expression regulation and demonstrated associations with various infertility-related factors, miRNAs represent promising candidates for diagnostic biomarker development.

The Role of MiRNAs in Epigenetics and Infertility

Epigenetics refers to changes in gene activity that do not involve changes in the underlying DNA sequence. These changes can be modulated by various environmental factors, such as diet status, psychological stress, and exposure to environmental toxins. Importantly, such epigenetic changes can significantly influence the expression of genes involved in reproductive health, which can affect fertility.

Research indicated that epigenetic modifications could contribute to both male and female infertility. In men, abnormal DNA methylation patterns in spermatozoa could affect sperm quality and motility and lead to problems with conception. In women, epigenetic alterations in ovarian or endometrial tissues might disrupt folliculogenesis, hormonal regulation, and embryo implantation success.

In addition, epigenetic changes can also be passed from one generation to another, potentially affecting the offspring's fertility through mechanisms collectively termed transgenerational epigenetic inheritance.^{7, 8} This phenomenon underscores the critical need to elucidate epigenetic contributions to infertility. Among epigenetic regulators, miRNAs are known to exert

epigenetic effects, altering gene expression and production of specific proteins.⁹ MiRNAs affect fertility by regulating various cellular processes, such as sperm production in men and the ovarian cycle in women.^{10, 11} Consequently, changes in miRNA levels can affect these processes, potentially leading to infertility.

By studying the epigenetic mechanisms involved in infertility, researchers hope to develop novel diagnostic modalities and targeted therapeutic strategies to enhance fertility outcomes for couples trying to conceive. The identification and modulation of epigenetic contributors to infertility could enable more precise, personalized treatment approaches. In addition, improved understanding of epigenetic influences on fertility might also lead to lifestyle modifications to optimize outcomes in assisted reproductive technologies (ART).

MicroRNAs in Female Infertility: Polycystic Ovary Syndrome (PCOS) and miRNAs

Polycystic ovary syndrome (PCOS) affects approximately 5-10% of reproductive-aged women, representing one of the most prevalent endocrine disorders in this population.¹² The diagnostic triad of PCOS includes chronic anovulation, polycystic ovarian morphology, and clinical and/or biochemical hyperandrogenism.^{13, 14} Importantly, PCOS demonstrates strong associations with metabolic disorders, such as type 2 diabetes (T2D), insulin resistance, and obesity.^{13, 14} While the precise etiology of PCOS remains unclear, current research implicates multifactorial influences including genetic predisposition, environmental factors, and iatrogenic effects such as chemotherapy exposure.¹⁵ Recent studies revealed distinct miRNA expression profiles in women with PCOS compared to healthy women, suggesting potential involvement of miRNAs in PCOS pathogenesis.¹⁶ This section aimed to evaluate the research in this area and discuss probable associations between miRNA dysregulation in PCOS, with particular focus on establishing mechanistic links that may inform novel therapeutic strategies. According to a study, PCOS patients had significantly higher levels of miRNAs, including miR-21, miR-155, miR-103, and miR-27b, than healthy women (table 1),¹⁷ which further supported the potential role of miRNAs in PCOS development and progression. Multiple studies demonstrated significant alterations in circulating miRNA profiles in women with PCOS. Specifically, serum levels of miR-16, miR-222, miR-106b, miR-19a, miR-146a, miR-30c, miR-186, and miR-24 were consistently increased in women with PCOS, while miR-320 was reduced compared to healthy

Table 1: Differential expressions of miRNAs in polycystic ovary syndrome and endometriosis

Differential expressions	PCOS	Endometriosis
Increase	miR-21, miR-103, miR-155, miR-27b, miR-222, miR-106b, miR-19a, miR-146a, miR-16, miR-30c, miR-24, miR-5706, miR-let-7i-3 pm, miR-4463, miR-638, miR-3665	miR-145, miR-99a, miR-143, miR-126, miR-99b, miR100, miR-150, miR125-b, miR-125a, miR-194, miR-223, miR-365, miR-1, miR-29c, miR-22-3p, miR-20a, miR-320a
Decrease	miR-320, miR-128, miR-124-3p, miR-let-7c, miR-29a-3p	miR-200a, miR-200b, miR-141, miR-424, miR-142-3p, miR-34c, miR-196b, miR-20a

PCOS: Polycystic ovary syndrome

controls (table 1).^{18, 19} Therefore, miR-146a and miR-30c exhibited particularly strong diagnostic potential for PCOS.²⁰ Increased expression of miR-222 was associated with type 2 diabetes and gestational diabetes pathogenesis.²¹

Additional studies revealed significant dysregulation of specific miRNAs in PCOS patients, with five miRNAs (let-7i-3 pm, miR-5706, miR-4463, miR-3665, and miR-638) showing increased expression and four miRNAs (miR-124-3p, miR-128, miR-29a-3p, and let-7c) demonstrating decreased expression compared to controls (table 1).²² Bioinformatics analyses of these altered miRNAs suggested that they participate in multiple pathological mechanisms underlying PCOS, including metabolic dysfunction, inflammatory responses, programmed cell death (apoptosis), vascular development (angiogenesis), immune system function, cellular energy metabolism (ATP binding), MAPK signaling cascades, and p53 pathways.^{16, 23} These consistent findings suggested that changes in miRNA levels might contribute to the PCOS development and progression.

Research has also explored the relationship between abnormal androgen production in PCOS and miRNA. In PCOS patients, free testosterone levels were strongly associated with miRNA levels, such as miR-21, miR-103, and miR-155, suggesting that aberrant miRNA expression might contribute to PCOS characteristics.²⁴

While miRNA research has advanced considerably, the precise mechanisms by which miRNAs contribute to PCOS pathophysiology remain under investigation. Profiling miRNAs can provide new insights into these molecular changes, offering therapeutic and prognostic information.²⁵ However, considerable controversy persists regarding optimal clinical strategies for symptom management, fertility improvement, and prevention of long-term PCOS complications.²⁶ A better understanding of these relationships and underlying processes could ultimately lead to more effective interventions and improved outcomes for individuals with PCOS.

Endometriosis and MiRNAs

Endometriosis is a gynecological disorder

characterized by the growth of endometrial tissue outside the uterine cavity. Emerging evidence suggested that miRNAs might serve as natural post-transcriptional regulators involved in the development of endometriotic lesions.²⁷ The current gold standard for definitive endometriosis diagnosis remains laparoscopic surgical evaluation, an invasive procedure with inherent patient risks.²⁸ Therefore, developing new non-invasive biomarkers to diagnose endometriosis in the early stages is an urgent need.²⁹ A previous study identified serum miRNAs as promising diagnostic biomarkers,²⁶ with 14 upregulated miRNAs (miR-145, miR-143, miR-99a, miR-99b, miR-126, miR100, miR125-b, miR-150, miR-125a, miR-223, miR-194, miR-365, miR-29c, and miR-1) and eight downregulated miRNAs (miR-200a, miR-141, miR-200b, miR-142-3p, miR-424, miR-34c, miR-20a, and miR-196b) identified (table 1).³⁰ In another study, researchers showed that in patients with endometriosis, the miR-9 and miR-34 families had abnormalities in the atopic endometrium.³¹ For example, studies showed elevated miR-20a levels in endometriosis lesions.³² The researchers suggested that dysregulation of these miRNAs and their target genes could contribute to the proliferative phenotype observed during the early secretory phase of the menstrual cycle in affected women.³³ As a result, many miRNAs have been identified as regulators of gene expression involved in inflammatory responses, local estrogen biosynthesis, progesterone resistance, cell invasion, extracellular matrix regeneration, and angiogenesis in endometriosis.³⁴ Six miRNAs (i.e., miR-134-5p, miR-197-5p, miR-22-3p, miR-320a, miR-494-3p, and miR-939-5p) were selected as biomarkers of endometriosis and validated using qRT-PCR (table 1).³⁵ Among these, miR-22-3p and miR-320a exhibited significantly increased expression in the serum exosomes of patients with endometriosis (table 1).³⁶ Considering that endometriosis diagnosis and treatment are frequently delayed due to nonspecific symptoms and the lack of reliable biomarkers, with laparoscopic surgery remaining the diagnostic gold standard, there remains a

Table 2: Differential expressions of miRNAs in male infertility

Differential expressions	Expressions of miRNAs in male infertility
Increase	miR-145, miR-302a, miR-491-3p, miR-122-5p
Decrease	miR-122, miR-449a, miR-23b, miR-383, miR-520d-3p, miR-34b, miR-34c, miR-34c-5p, miR-122, miR-146b-5p, miR-181a, miR-374b, miR-509-5p, and miR-513a-5p

critical need to identify effective non-invasive indicators for early stages of the disease.

MicroRNAs in Male Infertility

Male infertility represents a multifactorial reproductive disorder accounting for over 50% of couples' infertility cases.³⁷ This condition may arise from diverse etiologies, including genital tract infections, hormonal disorders, genitourinary tract infections, genetic mutations, and testicular hyperthermia. The latter may occur in clinical contexts such as varicocele or behavioral patterns including prolonged driving with a vehicle, or habitual laptop use on the lap, a documented risk factor for elevated scrotal temperature.³⁸ Currently, semen analysis serves as the standard diagnostic tool for male infertility. However, this method has some limitations and fails to identify the cause of infertility in approximately 40% of cases, highlighting the need for more effective diagnostic tools. Biomarkers represent one of the most promising emerging diagnostic tools. These molecules, detectable in body fluids, can indicate disease states such as infertility through abnormal expression patterns. Thus, miRNAs have emerged as potential biomarkers for different types of male infertility, including azoospermia, oligospermia, asthenospermia, and teratozoospermia.³⁸ In related research, scientists developed a transgenic mouse model in which the downregulation of miR-382-3p was prevented by overexpression in mature Sertoli cells (table 2).³⁹

Azoospermia, the most serious type of male infertility, affects 15-20% of infertile men and is classified into two main types: obstructive azoospermia (OA) and non-obstructive azoospermia (NOA).⁴⁰ NOA, resulting from impaired spermatogenesis, represents approximately 60% of azoospermia cases. In contrast, OA accounts for the remaining 40% and occurs when normally produced sperm cannot reach the ejaculate due to genital tract obstructions.⁴¹ NOA specifically refers to testicular failure to produce sperm and accounts for approximately 0.6% of all men, 10% of infertile men, and 60% of men with azoospermia.⁴¹ In recent years, the misplaced expression of miRNAs has been shown to affect several biological processes and was associated

with certain reproductive disorders in men, particularly those affecting spermatogenesis. Given the crucial role of miRNAs in all stages of spermatogenesis, their dysregulation logically contributes to numerous male infertility issues.⁴¹ A study showed decreased levels of miR-383 and miR-520d-3p and increased expression of miR-302a and miR-491-3p in NOA patients. It also reported that the expression of miR-17-92 and miR-371 was regulated in the testes of NOA patients (table 1).⁴² Another study found that decreased levels of miR-34b and miR-34c were associated with reduced male fertility, NOA, defective meiosis, impaired sperm maturation, and spermatogenesis disorders (table 2).⁴³

A similar study showed that dysregulation of miR-449 was associated with male infertility characterized by impaired sperm motility.⁴⁴ Comparative analysis of seminal plasma from fertile and infertile men revealed significantly reduced levels of seven miRNAs, including miR-34c-5p, miR-122, miR-146b-5p, miR-181a, miR-374b, miR-509-5p, and miR-513a-5p in azoospermic patients (table 2).⁴⁵ Another study found that miR-122-5p was upregulated in OA patients compared to NOA patients.⁴⁶⁻⁴⁸ These miRNAs could be used as diagnostic biomarkers for idiopathic male infertility and could inform the development of targeted therapies. Their irregularity could be the key to developing new and advanced diagnostic and prognostic approaches in male infertility. To take full advantage of their potential, further investigation is required to identify additional miRNAs involved in reproductive pathologies, the mechanism by which they work, and how their expression modulation might improve clinical outcomes. Collectively, these findings substantiate the regulatory role of miRNAs in male germ and somatic cells, where expression alterations could lead to reproductive disorders.

MiRNAs as Potential Biomarkers in Infertility Diagnosis

An ideal biomarker should be target-specific, stable, non-invasive, sensitive to detection, and easily accessible from accessible biological sources.⁴⁹ In clinical practice, serum proteins have conventionally served as biomarkers for disease detection and prognosis.⁵⁰ However, proteins as biomarkers have limitations, such

as reduced diagnostic value due to inadequate specificity and sensitivity. On the other hand, miRNAs emerge earlier in both intracellular and extracellular environments and can be detected with greater precision, enabling earlier disease identification. MiRNAs are typically expressed during specific biological phases.^{50, 51} Additionally, miRNAs are often observed as homogeneous populations, while proteins often undergo post-translational modifications and create heterogeneous forms.⁵²

Various diagnostic approaches have been developed for infertility assessment in both sexes, with biomarker-based methods representing the most recent advancement. Biomarkers can be utilized to detect diseases prior to clinical symptom manifestation. Among potential biomarkers, miRNAs are considered particularly promising due to their unique properties. Circulating (serum) miRNAs were initially employed as biomarkers for diagnosing B-cell lymphoma.⁵³ In another study, a decrease in miR-122, miR-181a, and miR-34c was observed in the serum of patients with oligoasthenospermia.^{54, 55} Several studies published in recent years demonstrated that miRNAs could be suitable biomarkers for diagnostic purposes.^{52, 56, 57}

Protein factors, DNA methylation, and non-coding RNAs are all present in the sperm and egg genomes.⁵⁸ During spermatogenesis and oogenesis in mammals, mRNAs undergo post-transcriptional and post-translational regulation in differentiated germ cells, enabling observation of miRNAs during both spermatogenesis and oocyte maturation.

MiRNAs and Infertility Treatment

The modification and manipulation of miRNA activity in the body is of great interest due to their abnormal expression and role in the pathogenesis of human diseases.⁵⁹ The use of anti-miRNA oligonucleotides to target disease-related miRNAs is the most widely used approach for understanding their *in vivo* function and shows promise for developing new miRNA-based therapies.^{60, 61} In addition to their potential use as biomarkers, miRNAs can be considered for therapeutic purposes in a personalized medical field related to male and female infertility.⁶² Researchers are exploring different methods to modulate miRNA levels for therapeutic purposes, including supplementation with microRNA mimics to either reduce or upregulate their expression, and using lab-engineered molecules called anti-miRNAs for conditions involving miRNA overexpression.⁶²

The therapeutic potential of miRNAs, along with advances in nanomedicine, could lead to the

development of novel fertility treatments for both male and female infertility.^{63, 64} With advances in genomics, researchers demonstrated that these small non-coding RNA molecules are essential for mammalian ovarian function. The most recent microRNA-based therapeutic strategies for infertility focus on addressing gonadal damage through multi-pathway targeting agents. Given existing treatment limitations, an integrated approach combining preventive measures with microRNA therapy represents both a practical and desirable clinical option.

Conclusion

Understanding the mechanisms that lead to infertility in both men and women is essential. Despite careful and expensive clinical examinations, including invasive techniques such as testicular biopsy, infertility sometimes remains undiagnosed. Current research has provided conclusive evidence on the role of miRNAs in regulating infertility. When combined with traditional diagnostic techniques, miRNAs have significantly improved infertility diagnosis. MiRNAs are affordable, readily available, and can be easily extracted from blood and semen, making them clinically practical. This diagnostic simplification may enable more effective treatment strategies after symptom and goal identification. To establish miRNAs as diagnostic biomarkers for various infertility types, developing standardized protocols and accurate diagnostic validation techniques would be essential.

Authors' Contribution

M.Kh: Study concept, data gathering, and drafting, L.N: Data gathering and drafting; MT.HG: Study concept, data gathering, and drafting, H.S; Data gathering and drafting, H.ZZ: Data gathering and drafting; A.S: Data gathering and drafting; N.GR: Data gathering and drafting; E.M: Data gathering and drafting; MA.S: Data gathering and drafting; All authors have read and approved the final manuscript and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Conflict of Interest: None declared.

References

- 1 Jimenez-Ortega RF, Ortega-Melendez AI, Patino N, Rivera-Paredes B, Hidalgo-Bravo

- A, Velazquez-Cruz R. The Involvement of microRNAs in Bone Remodeling Signaling Pathways and Their Role in the Development of Osteoporosis. *Biology (Basel)*. 2024;13. doi: 10.3390/biology13070505. PubMed PMID: 39056698; PubMed Central PMCID: PMCPCMC11273958.
- 2 Schwerk J, Jarret AP, Joslyn RC, Savan R. Landscape of post-transcriptional gene regulation during hepatitis C virus infection. *Curr Opin Virol*. 2015;12:75-84. doi: 10.1016/j.coviro.2015.02.006. PubMed PMID: 25890065; PubMed Central PMCID: PMCPCMC4472456.
- 3 Maziere P, Enright AJ. Prediction of microRNA targets. *Drug Discov Today*. 2007;12:452-8. doi: 10.1016/j.drudis.2007.04.002. PubMed PMID: 17532529.
- 4 Salas-Huetos A, James ER, Aston KI, Carrell DT, Jenkins TG, Yeste M. The role of miRNAs in male human reproduction: a systematic review. *Andrology*. 2020;8:7-26. doi: 10.1111/andr.12714. PubMed PMID: 31578810.
- 5 Sweeney MF, Hasan N, Soto AM, Sonnenschein C. Environmental endocrine disruptors: Effects on the human male reproductive system. *Rev Endocr Metab Disord*. 2015;16:341-57. doi: 10.1007/s11154-016-9337-4. PubMed PMID: 26847433; PubMed Central PMCID: PMCPCMC4803593.
- 6 Reza A, Choi YJ, Han SG, Song H, Park C, Hong K, et al. Roles of microRNAs in mammalian reproduction: from the commitment of germ cells to peri-implantation embryos. *Biol Rev Camb Philos Soc*. 2019;94:415-38. doi: 10.1111/brv.12459. PubMed PMID: 30151880; PubMed Central PMCID: PMCPCMC7379200.
- 7 Gunes S, Esteves SC. Role of genetics and epigenetics in male infertility. *Andrologia*. 2021;53:e13586. doi: 10.1111/and.13586. PubMed PMID: 32314821.
- 8 Bar-Sadeh B, Rudnizky S, Pnueli L, Bentley GR, Stoger R, Kaplan A, et al. Unravelling the role of epigenetics in reproductive adaptations to early-life environment. *Nat Rev Endocrinol*. 2020;16:519-33. doi: 10.1038/s41574-020-0370-8. PubMed PMID: 32620937.
- 9 Nalbant E, Akkaya-Ulum YZ. Exploring regulatory mechanisms on miRNAs and their implications in inflammation-related diseases. *Clin Exp Med*. 2024;24:142. doi: 10.1007/s10238-024-01334-y. PubMed PMID: 38958690; PubMed Central PMCID: PMCPCMC11222192.
- 10 Nadri P, Nadri T, Gholami D, Zahmatkesh A, Hosseini Ghaffari M, Savvulidi Vargova K, et al. Role of miRNAs in assisted reproductive technology. *Gene*. 2024;927:148703. doi: 10.1016/j.gene.2024.148703. PubMed PMID: 38885817.
- 11 Salmasi S, Heidar MS, Khaksary Mahabady M, Rashidi B, Mirzaei H. MicroRNAs, endometrial receptivity and molecular pathways. *Reprod Biol Endocrinol*. 2024;22:139. doi: 10.1186/s12958-024-01304-9. PubMed PMID: 39529197; PubMed Central PMCID: PMCPCMC11552404.
- 12 Naserpoor L, Jannatifar R, Roshanaei K, Khoshandam M, Kallhor N. Association of rs13429458 and rs12478601 Single Nucleotide Polymorphisms of THADA Gene with Polycystic Ovary Syndrome. *Int J Fertil Steril*. 2022;16:36-41. doi: 10.22074/IJFS.2021.524795.1090. PubMed PMID: 35103430; PubMed Central PMCID: PMCPCMC8808251.
- 13 Norman RJ, Dewailly D, Legro RS, Hickey TE. Polycystic ovary syndrome. *Lancet*. 2007;370:685-97. doi: 10.1016/S0140-6736(07)61345-2. PubMed PMID: 17720020.
- 14 Stener-Victorin E, Teede H, Norman RJ, Legro R, Goodarzi MO, Dokras A, et al. Polycystic ovary syndrome. *Nat Rev Dis Primers*. 2024;10:27. doi: 10.1038/s41572-024-00511-3. PubMed PMID: 38637590.
- 15 Sadeghi HM, Adeli I, Calina D, Docea AO, Mousavi T, Daniali M, et al. Polycystic Ovary Syndrome: A Comprehensive Review of Pathogenesis, Management, and Drug Repurposing. *Int J Mol Sci*. 2022;23. doi: 10.3390/ijms23020583. PubMed PMID: 35054768; PubMed Central PMCID: PMCPCMC8775814.
- 16 Chen Z, Ou H, Wu H, Wu P, Mo Z. Role of microRNA in the Pathogenesis of Polycystic Ovary Syndrome. *DNA Cell Biol*. 2019;38:754-62. doi: 10.1089/dna.2019.4622. PubMed PMID: 31305133.
- 17 Xia H, Zhao Y. miR-155 is high-expressed in polycystic ovarian syndrome and promotes cell proliferation and migration through targeting PDCD4 in KGN cells. *Artif Cells Nanomed Biotechnol*. 2020;48:197-205. doi: 10.1080/21691401.2019.1699826. PubMed PMID: 31851829.
- 18 Javadi M, Rad JS, Farashah MSG, Roshangar L. An Insight on the Role of Altered Function and Expression of Exosomes and MicroRNAs in Female Reproductive Diseases. *Reprod Sci*. 2022;29:1395-407. doi: 10.1007/s43032-021-00556-9. PubMed PMID: 33825167.
- 19 Hu J, Tang T, Zeng Z, Wu J, Tan X, Yan J. The expression of small RNAs in exosomes

- of follicular fluid altered in human polycystic ovarian syndrome. *PeerJ*. 2020;8:e8640. doi: 10.7717/peerj.8640. PubMed PMID: 32117643; PubMed Central PMCID: PMC7035867.
- 20 Ebrahimi SO, Reisi S, Parchami Barjui S. Increased risk of polycystic ovary syndrome (PCOS) associated with CC genotype of miR-146a gene variation. *Gynecol Endocrinol*. 2018;34:793-7. doi: 10.1080/09513590.2018.1460341. PubMed PMID: 29637801.
- 21 Filardi T, Catanzaro G, Grieco GE, Splendiani E, Trocchianesi S, Santangelo C, et al. Identification and Validation of miR-222-3p and miR-409-3p as Plasma Biomarkers in Gestational Diabetes Mellitus Sharing Validated Target Genes Involved in Metabolic Homeostasis. *Int J Mol Sci*. 2022;23. doi: 10.3390/ijms23084276. PubMed PMID: 35457094; PubMed Central PMCID: PMC9028517.
- 22 Montazerian M, Yasari F, Aghaalkhani N. Ovarian extracellular MicroRNAs as the potential non-invasive biomarkers: An update. *Biomed Pharmacother*. 2018;106:1633-40. doi: 10.1016/j.biopha.2018.07.073. PubMed PMID: 30119239.
- 23 Babayev E, Duncan FE. Age-associated changes in cumulus cells and follicular fluid: the local oocyte microenvironment as a determinant of gamete quality. *Biol Reprod*. 2022;106:351-65. doi: 10.1093/biolre/iaab241. PubMed PMID: 34982142; PubMed Central PMCID: PMC8862720.
- 24 Butler AE, Ramachandran V, Cunningham TK, David R, Gooderham NJ, Benurwar M, et al. Increased MicroRNA Levels in Women With Polycystic Ovarian Syndrome but Without Insulin Resistance: A Pilot Prospective Study. *Front Endocrinol (Lausanne)*. 2020;11:571357. doi: 10.3389/fendo.2020.571357. PubMed PMID: 33101204; PubMed Central PMCID: PMC7556216.
- 25 Zhang C, Yu C, Lin Z, Pan H, Li K, Ma H. MiRNAs expression profiling of rat ovaries displaying PCOS with insulin resistance. *Arch Gynecol Obstet*. 2020;302:1205-13. doi: 10.1007/s00404-020-05730-z. PubMed PMID: 32757043; PubMed Central PMCID: PMC7524818.
- 26 Zafari N, Bahramy A, Majidi Zolbin M, Emadi Allahyari S, Farazi E, Hassannejad Z, et al. microRNAs as novel diagnostic biomarkers in endometriosis patients: a systematic review and meta-analysis. *Expert Rev Mol Diagn*. 2022;22:479-95. doi: 10.1080/14737159.2021.1960508. PubMed PMID: 34304687.
- 27 Kiesel L, Sourouni M. Diagnosis of endometriosis in the 21st century. *Climacteric*. 2019;22:296-302. doi: 10.1080/13697137.2019.1578743. PubMed PMID: 30905186.
- 28 Restaino S, Mereu L, Finelli A, Spina MR, Marini G, Catena U, et al. Robotic surgery vs laparoscopic surgery in patients with diagnosis of endometriosis: a systematic review and meta-analysis. *J Robot Surg*. 2020;14:687-94. doi: 10.1007/s11701-020-01061-y. PubMed PMID: 32146573.
- 29 Giampaolino P, Foreste V, Della Corte L, Di Filippo C, Iorio G, Bifulco G. Role of biomarkers for early detection of ovarian cancer recurrence. *Gland Surg*. 2020;9:1102-11. doi: 10.21037/gs-20-544. PubMed PMID: 32953625; PubMed Central PMCID: PMC7475347.
- 30 Tamaru S, Kajihara T, Mizuno Y, Mizuno Y, Tochigi H, Ishihara O. Endometrial microRNAs and their aberrant expression patterns. *Med Mol Morphol*. 2020;53:131-40. doi: 10.1007/s00795-020-00252-8. PubMed PMID: 32350620.
- 31 Koninckx PR, Ussia A, Adamyan L, Gomel V, Martin DC. Peritoneal fluid progesterone and progesterone resistance in superficial endometriosis lesions. *Hum Reprod*. 2022;37:203-11. doi: 10.1093/humrep/deab258. PubMed PMID: 34849906.
- 32 Kolanska K, Bendifallah S, Canlorbe G, Mekinian A, Touboul C, Aractingi S, et al. Role of miRNAs in Normal Endometrium and in Endometrial Disorders: Comprehensive Review. *J Clin Med*. 2021;10. doi: 10.3390/jcm10163457. PubMed PMID: 34441754; PubMed Central PMCID: PMC8396961.
- 33 Loke H, Rainczuk K, Dimitriadis E. MicroRNA Biogenesis Machinery Is Dysregulated in the Endometrium of Infertile Women Suggesting a Role in Receptivity and Infertility. *J Histochem Cytochem*. 2019;67:589-99. doi: 10.1369/0022155419854064. PubMed PMID: 31145039; PubMed Central PMCID: PMC6669859.
- 34 Raja MHR, Farooqui N, Zuberi N, Ashraf M, Azhar A, Baig R, et al. Endometriosis, infertility and MicroRNA's: A review. *J Gynecol Obstet Hum Reprod*. 2021;50:102157. doi: 10.1016/j.jogoh.2021.102157. PubMed PMID: 33957270.
- 35 Kimber-Trojnar Z, Pilszyk A, Niebrzydowska M, Pilszyk Z, Ruszala M, Leszczynska-Gorzela B. The Potential of Non-Invasive Biomarkers for Early Diagnosis of Asymptomatic Patients with Endometriosis. *J Clin*

- Med. 2021;10. doi: 10.3390/jcm10132762. PubMed PMID: 34201813; PubMed Central PMCID: PMC8268879.
- 36 Wu Y, Yuan W, Ding H, Wu X. Serum exosomal miRNA from endometriosis patients correlates with disease severity. *Arch Gynecol Obstet.* 2022;305:117-27. doi: 10.1007/s00404-021-06227-z. PubMed PMID: 34542679; PubMed Central PMCID: PMC8782809.
 - 37 Fath-Bayati L, Naserpour L, Khoshandam M, Jannatifar R, Fazaeli H. Recent advances in developing 3D culture systems of spermatogonial stem cell preservation and differentiation: A narrative review. *Int J Reprod Biomed.* 2023;21:681-96. doi: 10.18502/ijrm.v21i9.14397. PubMed PMID: 37969562; PubMed Central PMCID: PMC810643686.
 - 38 Chereshevnev VA, Pichugova SV, Beikin YB, Cheresheva MV, Iukhta AI, Stroeve YI, et al. Pathogenesis of Autoimmune Male Infertility: Juxtacrine, Paracrine, and Endocrine Dysregulation. *Pathophysiology.* 2021;28:471-88. doi: 10.3390/pathophysiology28040030. PubMed PMID: 35366245; PubMed Central PMCID: PMC830464.
 - 39 Gupta A, Mandal K, Singh P, Sarkar R, Majumdar SS. Declining levels of miR-382-3p at puberty trigger the onset of spermatogenesis. *Mol Ther Nucleic Acids.* 2021;26:192-207. doi: 10.1016/j.omtn.2021.07.001. PubMed PMID: 34513304; PubMed Central PMCID: PMC8413679.
 - 40 Kasak L, Laan M. Monogenic causes of non-obstructive azoospermia: challenges, established knowledge, limitations and perspectives. *Hum Genet.* 2021;140:135-54. doi: 10.1007/s00439-020-02112-y. PubMed PMID: 31955275.
 - 41 Kuyucu Y, Coskun G, Saker D, Karaoglan O, Urunsak IF, Izol V, et al. Immunohistochemical examination of androgen receptor and estrogen receptor alpha expressions in obstructive and non-obstructive azoospermia. *Syst Biol Reprod Med.* 2021;67:463-70. doi: 10.1080/19396368.2021.1958094. PubMed PMID: 34420467.
 - 42 Chen Y, Li X, Liao H, Leung X, He J, Wang X, et al. CFTR mutation compromises spermatogenesis by enhancing miR-15b maturation and suppressing its regulatory target CDC25A. *Biol Reprod.* 2019;101:50-62. doi: 10.1093/biolre/iox062. PubMed PMID: 30985893.
 - 43 Zhang HT, Zhang Z, Hong K, Tang WH, Liu DF, Mao JM, et al. Altered microRNA profiles of testicular biopsies from patients with nonobstructive azoospermia. *Asian J Androl.* 2020;22:100-5. doi: 10.4103/aja.aja_35_19. PubMed PMID: 31134916; PubMed Central PMCID: PMC6958976.
 - 44 Pantos K, Grigoriadis S, Tomara P, Louka I, Maziotis E, Pantou A, et al. Investigating the Role of the microRNA-34/449 Family in Male Infertility: A Critical Analysis and Review of the Literature. *Front Endocrinol (Lausanne).* 2021;12:709943. doi: 10.3389/fendo.2021.709943. PubMed PMID: 34276570; PubMed Central PMCID: PMC8281345.
 - 45 Kiani M, Salehi M, Mogheiseh A. MicroRNA expression in infertile men: its alterations and effects. *Zygote.* 2019;27:263-71. doi: 10.1017/S0967199419000340. PubMed PMID: 31412971.
 - 46 Saebnia N, Neshati Z, Bahrami AR. Role of microRNAs in etiology of azoospermia and their application as non-invasive biomarkers in diagnosis of azoospermic patients. *J Gynecol Obstet Hum Reprod.* 2021;50:102207. doi: 10.1016/j.jogoh.2021.102207. PubMed PMID: 34407467.
 - 47 Daneshmandpour Y, Bahmanpour Z, Hamzeiy H, Mazaheri Moghaddam M, Mazaheri Moghaddam M, Khademi B, et al. MicroRNAs association with azoospermia, oligospermia, asthenozoospermia, and teratozoospermia: a systematic review. *J Assist Reprod Genet.* 2020;37:763-75. doi: 10.1007/s10815-019-01674-9. PubMed PMID: 32189180; PubMed Central PMCID: PMC7183035.
 - 48 Cannarella R, Barbagallo F, Crafa A, La Vignera S, Condorelli RA, Calogero AE. Seminal Plasma Transcriptome and Proteome: Towards a Molecular Approach in the Diagnosis of Idiopathic Male Infertility. *Int J Mol Sci.* 2020;21. doi: 10.3390/ijms21197308. PubMed PMID: 33022946; PubMed Central PMCID: PMC7582752.
 - 49 Bellassai N, D'Agata R, Jungbluth V, Spoto G. Surface Plasmon Resonance for Biomarker Detection: Advances in Non-invasive Cancer Diagnosis. *Front Chem.* 2019;7:570. doi: 10.3389/fchem.2019.00570. PubMed PMID: 31448267; PubMed Central PMCID: PMC6695566.
 - 50 Condrat CE, Thompson DC, Barbu MG, Bugnar OL, Boboc A, Cretoiu D, et al. miRNAs as Biomarkers in Disease: Latest Findings Regarding Their Role in Diagnosis and Prognosis. *Cells.* 2020;9. doi: 10.3390/cells9020276. PubMed PMID: 31979244; PubMed Central PMCID: PMC7072450.
 - 51 Gareev I, Beylerli O, Zhao B. MiRNAs as

- potential therapeutic targets and biomarkers for non-traumatic intracerebral hemorrhage. *Biomark Res.* 2024;12:17. doi: 10.1186/s40364-024-00568-y. PubMed PMID: 38308370; PubMed Central PMCID: PMC810835919.
- 52 Velazquez-Cruz A, Banos-Jaime B, Diaz-Quintana A, De la Rosa MA, Diaz-Moreno I. Post-translational Control of RNA-Binding Proteins and Disease-Related Dysregulation. *Front Mol Biosci.* 2021;8:658852. doi: 10.3389/fmolb.2021.658852. PubMed PMID: 33987205; PubMed Central PMCID: PMC8111222.
- 53 Drees EEE, Pegtel DM. Circulating miRNAs as Biomarkers in Aggressive B Cell Lymphomas. *Trends Cancer.* 2020;6:910-23. doi: 10.1016/j.trecan.2020.06.003. PubMed PMID: 32660885.
- 54 Munawar M, Liaqat I, Ali S. MicroRNAs and male infertility. *Recent Advances in Noncoding RNAs.* 2022;193. doi: 10.5772/intechopen.106757.
- 55 Bjorkgren I, Gylling H, Turunen H, Huhtaniemi I, Strauss L, Poutanen M, et al. Imbalanced lipid homeostasis in the conditional Dicer1 knockout mouse epididymis causes instability of the sperm membrane. *FASEB J.* 2015;29:433-42. doi: 10.1096/fj.14-259382. PubMed PMID: 25366345.
- 56 Grossman H, Shalgi R. A Role of MicroRNAs in Cell Differentiation During Gonad Development. *Results Probl Cell Differ.* 2016;58:309-36. doi: 10.1007/978-3-319-31973-5_12. PubMed PMID: 27300184.
- 57 Dutta S, Sengupta P, Mottola F, Das S, Husain A, Ashour A, et al. Crosstalk Between Oxidative Stress and Epigenetics: Unveiling New Biomarkers in Human Infertility. *Cells.* 2024;13. doi: 10.3390/cells13221846. PubMed PMID: 39594595; PubMed Central PMCID: PMC811593296.
- 58 Cheuqueman C, Maldonado R. Non-coding RNAs and chromatin: key epigenetic factors from spermatogenesis to transgenerational inheritance. *Biol Res.* 2021;54:41. doi: 10.1186/s40659-021-00364-0. PubMed PMID: 34930477; PubMed Central PMCID: PMC810866607.
- 59 Deilulis JA. MicroRNAs as regulators of metabolic disease: pathophysiologic significance and emerging role as biomarkers and therapeutics. *Int J Obes (Lond).* 2016;40:88-101. doi: 10.1038/ijo.2015.170. PubMed PMID: 26311337; PubMed Central PMCID: PMC810472234.
- 60 Walayat A, Yang M, Xiao D. Therapeutic Implication of miRNA in Human Disease. *Antisense therapy.* 2019;93. doi: 10.5772/intechopen.82738.
- 61 Ratti M, Lampis A, Ghidini M, Salati M, Mirchev MB, Valeri N, et al. MicroRNAs (miRNAs) and Long Non-Coding RNAs (lncRNAs) as New Tools for Cancer Therapy: First Steps from Bench to Bed-side. *Target Oncol.* 2020;15:261-78. doi: 10.1007/s11523-020-00717-x. PubMed PMID: 32451752; PubMed Central PMCID: PMC8107283209.
- 62 Vashisht A, Gahlay GK. Using miRNAs as diagnostic biomarkers for male infertility: opportunities and challenges. *Mol Hum Reprod.* 2020;26:199-214. doi: 10.1093/molehr/gaaa016. PubMed PMID: 32084276.
- 63 Alexandri C, Daniel A, Bruylants G, Demeestere I. The role of microRNAs in ovarian function and the transition toward novel therapeutic strategies in fertility preservation: from bench to future clinical application. *Hum Reprod Update.* 2020;26:174-96. doi: 10.1093/humupd/dmz039. PubMed PMID: 32074269.
- 64 Falagan-Lotsch P, Grzincic EM, Murphy CJ. New Advances in Nanotechnology-Based Diagnosis and Therapeutics for Breast Cancer: An Assessment of Active-Targeting Inorganic Nanoplateforms. *Bioconjug Chem.* 2017;28:135-52. doi: 10.1021/acs.bioconjchem.6b00591. PubMed PMID: 27973767.