

Exploring the Adverse Effects of Fenugreek in Humans: A Scoping Review

Ehsan Amiri Ardekani¹, PharmD, PhD;¹⁵
Mohammad Mahdi Parvizi^{2,3,4,5}, MD, PhD,
MPH;¹⁶ Roghaye Zare⁶, MD, PhD; Maryam
Mosaffa Jahromi³, PharmD, PhD; Thomas
Rampp⁷, MD, PhD; Mehdi Pasalar³, MD, PhD

¹Department of Phytopharmaceuticals
(Traditional Pharmacy), School of
Pharmacy, Shiraz University of Medical
Sciences, Shiraz, Iran;

²Molecular Dermatology Research Center,
Shiraz University of Medical Sciences,
Shiraz, Iran;

³Research Center for Traditional Medicine
and History of Medicine, Shiraz University
of Medical Sciences, Shiraz, Iran;

⁴Smart University of Medical Sciences,
Tehran, Iran;

⁵Department of Medical Journalism,
School of Paramedical Sciences,
Shiraz University of Medical Sciences,
Shiraz, Iran;

⁶Department of Persian Medicine, School
of Persian Medicine, Shahid Sadoughi
University of Medical Sciences, Ardakan,
Yazd, Iran;

⁷Department of Internal and Integrative
Medicine, Evang. Kliniken Essen-
Mitte, Faculty of Medicine, University of
Duisburg-Essen, Essen, Germany

Correspondence:

Mohammad Mahdi Parvizi, MD, PhD, MPH;
Molecular Dermatology Research Center,
Shahid Faghihi Hospital,
Postal code: 7134844119, Zand Blvd.,
Shiraz, Iran

Tel: +98 71 32125592

Email: mmparvizi@gmail.com

Received: 01 March 2025

Revised: 02 May 2025

Accepted: 28 June 2025

What's Known

- Fenugreek (*Trigonella foenum-graecum*) is a plant that has been widely used both as a culinary spice and a medicinal herb.

What's New

- The most frequently reported adverse effect associated with oral fenugreek ingestion is mild gastrointestinal discomfort.
- Other documented adverse events include hypoglycemia, suspected hypokalemia, allergic reactions, a maple syrup-like odor in the sweat or urine of mothers and breastfed infants, and interactions with specific medications.

Abstract

Background: *Trigonella foenum-graecum*, commonly known as fenugreek, is used both as a spice and a medicinal herb. While numerous studies investigated its therapeutic effects, this scoping review aimed to explore the reported adverse effects associated with fenugreek consumption in humans.

Methods: A systematic search of several scientific databases was conducted, including Google Scholar, Web of Science, PubMed/PMC-MEDLINE, Scopus, and Science Direct, from January 1990 to September 2024. The search utilized keywords such as “Fenugreek”, “Adverse Effects”, “Clinical Trial”, and “Case Report”, or “Case Series”. References of retrieved articles were also screened.

Results: The review included 60 articles. Of these, 14 clinical trials reported adverse effects, 23 reported none, and 13 did not provide information on adverse effects. Additionally, 10 case reports or case series (reported in eight articles) detailed allergic reactions or hypersensitivity signs and symptoms. The most common adverse effect was mild gastrointestinal discomfort following oral consumption. Other reported effects included hypoglycemia, potential hypokalemia, allergic reactions, a maple syrup odor in the urine, sweat, or skin of infants and mothers, and interactions with certain medications.

Conclusion: Fenugreek is generally considered safe, with most reported side effects being mild and self-limiting. No fatalities have been attributed to its use. This evidence might be valuable for both the general public and healthcare professionals.

Please cite this article as: Amiri Ardekani E, Parvizi MM, Zare R, Mosaffa Jahromi M, Rampp T, Pasalar M. Exploring the Adverse Effects of Fenugreek in Humans: A Scoping Review. Iran J Med Sci. doi: 10.30476/ijms.2025.106073.4017.

Keywords • Trigonella • Spice • Drug-related side effects and adverse reactions • Clinical trial • Persian medicine

Introduction

In recent years, public interest in complementary and alternative medicine (CAM) has grown significantly. Concurrently, global research publications on the application of CAM in clinical practice have increased. The use of herbal medicines and medicinal plants for common illnesses is becoming more popular, aligning with the rising integration of CAM alongside modern conventional treatments for a wide range of conditions.¹⁻⁴

This growing attention from both patients and the scientific community is partly due to the incomplete success of conventional medications in treating or managing numerous common ailments, including gynecological disorders, type 2 diabetes mellitus, male reproductive problems, hyperlipidemia, cardiovascular conditions, and COVID-19.⁵⁻¹⁰ Furthermore, several studies

documented a patient perception that the herbal and natural remedies are safer than synthetic pharmaceutical alternatives.⁵⁻¹⁰

Fenugreek (*Trigonella foenum-graecum* L.) is an annual plant from the *Fabaceae* family, native to some regions of Asia, Africa, southern Europe, and also Canada. The plant typically reaches 30-60 cm in height and features trifoliate green leaves. Its flowers produce slender, boat-shaped pods, each containing an average of 10-20 green-brown seeds.¹¹ Fenugreek seeds and leaves are used variously as a spice, food additive, flavoring, condiment, preservative, and vegetable.¹² For example, the seeds and leaves are utilized in diverse culinary applications, including Iranian stews, Swiss cheese flavoring, German syrup and bitter rum, and Egyptian mixed seed powder for flatbread. They are also used in curries, as dyes, and—when roasted—as a coffee substitute in Africa, while young seedlings are consumed as a vegetable.^{11, 13}

The pharmacotherapeutic potential of fenugreek is derived from its rich array of bioactive components, including steroids, polyphenols, alkaloids, saponins, hydrocarbons, and galactomannan fiber. Furthermore, the plant contains substantial quantities of furostanolic saponins, such as trigoneoside, isoorientin, vitexin, and isovitexin. The seeds are a source of triglycerides, fatty acids, polysaccharides, notably a high concentration of galactomannan and flavone C-glycosides.¹⁴⁻¹⁷ Nutritionally, fenugreek is rich in vitamin A, B₁, B₂, C, niacin, and nicotinic acid.¹⁸ The seeds contain 28.4% protein, 9.3% fiber, and 7.1% fat,¹⁹ in addition to minerals such as magnesium, iron, copper, chromium, and calcium.²⁰

Fenugreek seed oil contains omega-6 fatty acids, which are beneficial in managing coronary heart disease, inflammation, and cancer. It also contains palmitic acid, pinene, and other components with antioxidant activity.²¹ The oligosaccharides in fenugreek seeds are known to confer health benefits in type 2 diabetes by preventing the rapid absorption of monosaccharides,¹⁷ and fenugreek supplementation has been shown to significantly improve both fasting and postprandial blood glucose levels.²² Furthermore, diosgenin, a compound found in fenugreek, seems to prevent colon cancer²³ and has beneficial effects in hyperlipidemic patients.²⁴ Fenugreek extract was also demonstrated to increase serum total testosterone levels in males.²⁵ A saponin-rich fenugreek extract can generate multi-bioactive extracts that inhibit pancreatic lipase and cholesterol bioaccessibility, potentially leading to a hypocholesterolemic effect.²⁶ Additionally,

fenugreek seed is recognized for its ability to improve breast milk production.²⁷

Given the increasing popularity of using herbal medicine to treat various diseases and conditions,^{5, 9, 28, 29} there is a growing need to evaluate and report the adverse effects of medicinal plants such as fenugreek. Therefore, this scoping review aimed to comprehensively examine human studies on fenugreek consumption, including clinical trials, case reports, and case series, to identify any associated adverse effects. The findings of this study could provide valuable information for a wide audience, particularly for patients who use fenugreek as an herbal remedy.

Materials and Methods

Study Design

This scoping review aimed to collect information on the adverse effects associated with fenugreeks consumption in human studies. A systematic search was conducted using various keywords and related MeSH terms, such as “Trigonellas,” “*Trigonella foenum-graecum*,” “*Trigonella foenum graecum*,” “Fenugreek,” “Fenugreeks,” “Foenumgraecum,” “Adverse Effects,” “Clinical Trial”, and “Case Report” and “Case Series” in multiple databases and search engines, such as Google Scholar, Web of Science, PubMed, Scopus, and Science Direct, from January 1990 to September 2024. The reference lists of retrieved articles were also screened.

The study included only English-published articles that documented the adverse effects of fenugreeks on humans. Animal studies, non-English publications, and articles for which the full text was unavailable were excluded. The eligible articles were reviewed, and relevant data were extracted and recorded in Microsoft Excel software (Microsoft Press, Redmond, WA, USA). Finally, the included articles were categorized as clinical trials, case reports, or case series.

Data Extraction

For clinical trials, the extracted data included: the health status of the participants, the number of subjects in the fenugreek and control groups, the formulation, dosage, and administration schedule of fenugreek, the interval between consumption and follow-up, concomitant interventions, study design, and any reported adverse effects. For case reports and case series, the extracted information included article title, patient demographics (age, sex), health status, medical history, clinical examination

findings, as well as the formulation, dosage, administration schedule, and reported adverse effects of fenugreek. Following the review and data extraction, the studies were categorized, and the adverse effects observed in the selected studies were summarized using the available data and tables.

The study protocol was approved by the Research Ethics Committee of Shiraz University of Medical Sciences (code: IR.SUMS.MED.REC.1399.420).

Results

Based on the inclusion criteria, the relevant data from the selected articles were organized into tables. The final review included 60 articles, comprising 50 clinical trials and 10 case reports.

Clinical Trial Articles

Of the 50 clinical trials, 14 reported adverse effects associated with fenugreek consumption, 23 reported no adverse effects, and the remaining 13 did not mention adverse effects.

As shown in table 1, among the 14 clinical trials reporting adverse effects, 57 out of 384 participants in the fenugreek group experienced them. These studies involved healthy individuals,³⁰ women with low libido,³¹ lactating women,³² overweight patients,³³ and individuals with type 1 and type 2 diabetes.^{34, 35}

These reported adverse effects included gastrointestinal complications, such as reflux, abdominal pain, diarrhea, mild abdominal distention, hunger, appetite loss, and nausea.

Other complications included migraine exacerbation, maple syrup odor in urine or sweat, frequent urination, dizziness, increased need for sanitary napkins, and hypoglycemia. The most prevalent adverse effect was urine with a maple syrup odor.

The treatment duration varied across studies, ranging from a single dose of fenugreek leaf aqueous extract to a 12-week regimen of fenugreek seed capsules.

As shown in table 2, 23 clinical trial articles reported no adverse effects. The majority of these studies investigated type 2 diabetes and utilized fenugreek seed powder.

Research on the effects of fenugreek has been conducted across diverse populations. These included studies on individuals with dysmenorrhea (n=3), polycystic ovary syndrome (n=2), Parkinson's disease (n=1), pre-diabetes (n=1), and testosterone deficiency (n=1), as well as on healthy men (n=2) and mechanically ventilated patients (n=1). The duration of these interventions ranged from 3 days to 6 months.

As shown in table 3, 13 clinical trials on fenugreek consumption, involving over 500 participants, reported no adverse effects. The studies investigated a range of health conditions, including type 2 diabetes, obesity, menopausal symptoms, hernia surgery, lactation, low libido in women, and gingivitis.

The duration of these studies varied, with the shortest being a two-day investigation using a 10% fenugreek transdermal patch on hernia surgery wounds. The longest was a 6-month trial involving patients with type 2 diabetes.

Table 1: Adverse effects associated with *Trigonella foenum-graecum* (fenugreek) in clinical trials

Authors/ year of publication	Country	Participants	Health status	Preparation and dosage	Duration of treatment	Adverse effects reported from the fenugreek group	Adverse effects reported in the control group
Steel et al., 2011 ³⁶	Australia	30	Healthy men	Testophen tablets contain 300 mg of fenugreek powder two times a day	6 weeks	Three cases of mild stomach pain before meals	Was not observed
Rao et al., 2015 ³⁷	Australia	40	Women with low libido	600 mg of fenugreek seed extract per day	In two consecutive menstrual periods	Two cases of exacerbation of migraine, two cases of reflux	Was not observed
Rao et al., 2020 ³¹	Australia	50	Benign prostatic hyperplasia	600 mg of fenugreek seed extract per day	12 weeks	Three cases of reflux	One case of reflux
Najdi et al., 2019 ³⁸	Saudi Arabia	5	Type 2 diabetes that was treated with metformin	2 g of fenugreek in capsule form (one 500 mL capsule after breakfast, two capsules after lunch, and one capsule after dinner)	12 weeks	One case of hypoglycemia	Was not observed

Authors/ year of publication	Country	Participants	Health status	Preparation and dosage	Duration of treatment	Adverse effects reported from the fenugreek group	Adverse effects reported in the control group
Bumrungpert et al., 2018 ³²	Thailand	25	Lactating lady	3 capsules (200 mg fenugreek seeds, 120 mg ginger, 100 mg turmeric) 3 times a day	4 weeks	Two cases of flatulence, two cases of urine with the odor of maple syrup	Two cases of urine with the odor of maple syrup
Chevassus et al., 2009 ³³	France	24	Healthy	Fenugreek tablets (one group, 588 mg, and another group, 1176 mg)	3-14 days with 14 days in between	One case of heartburn, two cases of the specific smell of urine	Was not observed
Chevassus et al., 2010 ³⁹	France	19	Overweight	Fenugreek tablets contain 1176 mg of hydroalcoholic extract of fenugreek seeds daily	6 weeks	Four cases of mild gastrointestinal symptoms, one case of a specific odor of urine and sweat	Was not observed
Emtiazzy et al., 2018 ⁴⁰	Iran	28	Mild asthma	Ten mL of fenugreek seed extract syrup, two times a day	4 weeks	Two cases of increased sanitary napkin use during menstruation	Was not observed
Sharma et al., 1990 ³⁴	India	10	Type 1 diabetes	50 g of fenugreek seed powder, two times a day, at lunch and dinner, added to bread	10 days	Four cases of gastrointestinal symptoms, including diarrhea and bloating	Was not observed
Gupta et al., 2001 ³⁵	India	12	Type 2 diabetes	1 g of hydroalcoholic extract of fenugreek seeds in capsule form	2 months	Five people with mild abdominal distension	Was not observed
Abdel-Barry et al., 2000 ³⁰	Iraq	20	Healthy	40 mg per Kg bodyweight of fenugreek leaf extract in 10 mL of boiled water	A single dose	Four cases felt hungry, four cases of frequent urination, and four cases of dizziness.	Was not observed
Lu et al., 2008 ⁴¹	China	46	Type 2 diabetes	Six 35% pills from seeds three times a day	12 weeks	Two cases of nausea and one case of diarrhea	Was not observed
Sudheeran et al., 2016 ⁴²	India	20	Healthy	500 mg capsules, including 300 mg of fiber and 200 mg of turmeric, two times daily before breakfast and at bedtime	30 days	Two cases of decreased appetite	One case of gastro- intestinal problems
Mirgoaybayat et al., ⁴³	Iran	55	Polycystic Ovary Syndrome	Take 333 mg of fenugreek capsules three times a day (Mootta capsules contain 333 mg of dry fenugreek extract, standardized to contain 53.7% trigonelline, the key phytochemical compound in fenugreek)	2 months	Six (10.9%) cases of nausea	32 (58.2%) cases of nausea, 16 (29.1) cases of headache

Table 2: Clinical trials with no associated adverse effects from *Trigonella foenum-graecum* (fenugreek) administration

Authors/ year of publication	Country	Number of participants in the fenugreek group	Health status of the fenugreek group	Preparation and dosage	Duration of the treatment
Shamshad et al., 2016 ⁴⁴	India	44	Involved in menopausal symptoms	250 mg capsules of fenugreek seed extract twice a day for a week and four times a day for 12 weeks	90 days
Swaroop et al., 2015 ⁴⁵	USA	50	Polycystic ovary	500 mg capsules of fenugreek seed extract two times a day	90 days
Nathan et al., 2013 ⁴⁶	India	25	Parkinson	300 mg of fenugreek seed extract two times a day	6 months
Florentin et al., 2019 ⁴⁷	Greece	50	Pre-diabetes	Tablets containing bergamot extract 500 mg, fenugreek seed extract 200 mg, and olive leaf extract 100 mg once a day	6 months
Verma et al., 2016 ⁴⁸	India	77	Type 2 diabetes	500 mg capsules of fenugreek seed extract two times a day	90 days
Geberemeskel et al., 2019 ⁴⁹	Ethiopia	57	Type 2 diabetes	25 mg of fenugreek seed extract solution two times a day	1 month
Madar et al., 1988 ⁵⁰	Israel	21	Type 2 diabetes	15 g of fenugreek seeds mixed with water	7 days
Younesy et al., 2014 ⁵¹	Iran	51	Dysmenorrhea	900 mg capsules of fenugreek seed powder three times a day	The first three days of menstruation
Losso et al., 2009 ⁵²	USA	10	Type 2 diabetes	9% of bread wheat flour replaced with 2.5 g of fenugreek seed powder (1 slice, two times a day)	1 week
Inanmdar et al., 2016 ⁵³	India	20	Primary dysmenorrhea	Three capsules equivalent to 3 g of fenugreek seeds	The first 3 days of menstruation
Park et al., 2018 ⁵⁴	Korea	44	Testosterone deficiency syndrome	200 mg capsules of fenugreek seeds two times a day	8 weeks
Hassanzadeh et al., 2013 ⁵⁵	Iran	23	Polycystic ovary	500 mg capsules of fenugreek seed extract two times a day	2 months
Rao et al., 2016 ⁵⁶	Australia	55	Healthy men	600 mg daily fenugreek seed extract	12 weeks
Maheshwari et al., 2017 ⁵⁷	India	50	Healthy men	500 mg capsules of fenugreek seeds after breakfast	12 weeks
Rao et al., 2020 ³¹	Australia	40	Type 2 diabetes mellitus	Used two chapatis twice a day, 6 days/week for a daily dose of 5.45 g of an <i>Nigella sativa</i> /fenugreek combination	12 weeks
Hausenblas et al., 2021 ⁵⁸	USA	19	Healthy men	Fenugreek 400 mg/d	60 days
Zarghi et al., 2021 ⁵⁹	Iran	33	Mechanically ventilated patients hospitalized	3 mg of fenugreek seed powder with a gavage solution twice daily	5 days
Foroumandi et al., 2023 ⁶⁰	Iran	41	Alzheimer's disease	Received 5 mL oral seed extract of fenugreek (equivalent to 500 mg of dry extract) added to the similar routine treatment, including Donepezil (5 mg twice a day) and Sertraline (50 mg once a day).	4 months
Gupta et al., 2024 ¹⁶	India	42	Type 2 diabetes	Use 1000 mg (500 mg×2) daily (Fenfuro®) capsules, that was a novel Fenugreek seed extract with >45% furostanolic saponins	12 weeks
Lee-Ødegård et al., 2024 ⁶¹	Norway	600 mg (n=21), 1200 mg (n=25) and 1800 mg (n=27)	Men with reduced energy and libido related to non-optimal testosterone levels	Taking 3 tablets daily, 600 mg (n=21), 1200 mg (n=25), and 1800 mg (n=27) of fenugreek extract and essential nutrients. ^a	12 weeks
Hota D et al., 2024 ^{16, 62}	India	204 (total patients)	Type 2 diabetes	Fenfuro® in the dosage of 500 mg twice daily along with metformin	12 weeks
Singh et al., 2023 ⁶³	India	113	Pre-menopausal women with polycystic ovary syndrome (PCOS)	Furocyst® (2 capsules of 500mg/day)	90 days

a. No reported side effects, but with a slight increase in serum concentrations of ALAT and creatinine

Table 3: Clinical trials of *Trigonella foenum-graecum* (fenugreek), not reporting adverse effects

Authors/ year of publication	Country	Fenugreek group participant	Health status of the fenugreek group	Preparation and dosage	Duration of the treatment
Robert et al., 2016 ⁶⁴	Malaysia	10	Healthy	Buns and flatbreads each contain 10% fenugreek seed powder with 50 g of glucose and 250 mL of water daily.	6 times
Sundaram et al., 2018 ⁶⁵	India	40	Uncontrolled type 2 diabetes treated with metformin, with chronic periodontal disease	12.5 g of fenugreek powder before breakfast and lunch	1 month
Ansari et al., 2019 ⁶⁶	Iran	30	After transdermal local patch hernia surgery	10% fenugreek daily	2 days
Bordia et al., 1997 ⁶⁷	India	20	Non-insulin-dependent diabetes mellitus (NIDDM) and coronary artery disease	2.5 g twice a day	3 months
	India	20	Non-insulin-dependent diabetes mellitus (NIDDM) without coronary artery disease	2.5 g twice a day	3 months
	India	20	Healthy	2.5 g twice a day	3 months
Ghasemi et al., 2015 ²⁷	Iran	39	Healthy breastfeeding mothers	7.5 g of fenugreek seed powder with 3 g of black tea three times a day	4 weeks
Kassaian et al., 2009 ⁶⁸	Iran	11	Type 2 diabetes	10 g of fenugreek seed powder soaked in water daily	8 weeks
Kassaian et al., 2009 ⁶⁸	Iran	7	Type 2 diabetes	10 g of fenugreek seed powder in yogurt daily	8 weeks
Ranad et al., 2017 ⁶⁹	India	30	Type 2 diabetes	10 g of fenugreek seeds soaked in water	6 months
Kiss et al., 2018 ⁷⁰	Hungary	8	Healthy	500 mg capsules, the first day two capsules in two meals at noon and evening, the last day two capsules in the evening, and on other days two capsules three times a day	11 days
Mathern et al., 2009 ⁷¹	USA	18	Fat	Extracts of 4 and 8 g of fenugreek fiber in beer	3 days
Steels et al., 2017 ⁷²	Australia	54	Menopausal symptoms	600 mg of seed extract without fenugreek pod daily	12 weeks
Palacios et al., 2019 ⁷³	Spain	29	Women with low libido	Tablets contain 300 mg of fenugreek extract two times a day	2 months
Mehrzadi et al., 2020 ⁷⁴	Iran	150	Type 2 diabetes	Traditional herbal capsules including 115 mg fenugreek, caper, rosehip, <i>Securigera securidaca</i> , <i>Silybum marianum</i> (milk thistle), nettle, and <i>Caucasian whortleberry</i> daily	3 months
Varghese et al., 2021 ⁷⁵	Karnataka	15	Gingivitis	Use toothpaste daily, two times (morning and at night) in a pea-size amount	14 days

Case Reports and Case Series Articles

As detailed in table 4, ten case reports were qualified, reviewed, and analyzed. The cases involved seven females and three males, with ages ranging from infancy to 67 years.

The most commonly reported adverse effects of fenugreek were allergic reactions, including pruritus, shortness of breath, rhinorrhea, diarrhea, angioedema, cough, and anaphylaxis (seven cases). Almost all cases reported a history of allergies to other substances, such as peas, peanuts, and coriander. In one case, the topical application of a fenugreek ointment to the scalp for dandruff resulted in syncope.

One case of Stevens-Johnson syndrome and toxic epidermal necrosis was reported in a 32-year-old woman. The reaction, presenting

as blisters and sores on her face and upper torso, occurred 1 month postpartum after she had taken a fenugreek-containing medication to stimulate lactation.

In another report, a newborn boy exhibited a maple syrup odor from his skin and diaper area after his mother had consumed fenugreek during early labor.

Another case described a 67-year-old woman with atrial fibrillation whose international normalized ratio (INR) increased following the consumption of fenugreek and Boldo drops while on warfarin therapy.

In another report, symptoms suggestive of serotonin syndrome, including hyperreflexia, tachycardia, nausea, and anxiety, were observed in a 38-year-old woman with psychosis.

Table 4: Reported adverse events of *Trigonella foenum-graecum* (fenugreek) in case reports and series

Authors/ year of publication	Country	Sex	Age	Health status	How to expose	Adverse effects	Result
Patil et al., 1997 ⁷⁶	India	Woman	37	Mild asthma - pea allergy	Open and smell the glass containing fenugreek	Runny nose and eyes, cough, fainting	Treated
Patil et al., 1997 ⁷⁶	India	Woman	45	Asthma - Dandruff with a history of wheezing after eating fenugreek	Fenugreek seed ointment on the scalp	Angioedema of the face, runny nose, and numbness of the head	Treated
Bentele- Jaberg et al., 2015 ⁷⁷	Switzerland	Woman	32	1 month after delivery	Eat medicine made from fenugreek seeds	Blisters and sores on the face and upper body, with involvement of the mouth, lips, and tongue	Treated
Joseph et al., 2018 ⁷⁸	USA	Man	14	Healthy	Eat food containing fenugreek	Urticaria, chest tightness, abdominal pain, vomiting	Treated
Korman et al., 2001 ⁷⁹	Israel	Man	Infant	Healthy	The baby's mother consumed fenugreek in the first hours of labor pains	The smell of maple syrup from the skin and diaper area	Treated
Lambert et al., 2001 ⁸⁰	Canada	Woman	67	Atrial fibrillation on warfarin	A fenugreek capsule with ten drops of Boldo after a meal	Increase INR	Treated
Aurich et al., 2019 ⁸¹	Germany	Woman	34	Atopic dermatitis, asthma, peanut allergy	Eat a Chinese soup containing fenugreek	Facial flushing, angioedema, shortness of breath, nausea, and diarrhea	Treated
Doolabh et al., 2019 ⁸²	Australia	Woman	38	Psychosis on sertraline during breastfeeding	Use fenugreek supplements to increase breastfeeding	Symptoms of serotonin syndrome include hyperreflexia, nausea, anxiety, and tachycardia	Treated
Ohnuma et al., 1998 ⁸³	Japan	Woman	26	Healthy	Use curry powder containing fenugreek	Itching, diarrhea, wheezing	Treated
Ebo et al., 2006 ⁸⁴	Belgium	Man	25	Healthy with a history of allergies to fenugreek and coriander in occupational exposure	Eat a loaf of bread containing fenugreek and coriander	Anaphylactic reactions include generalized urticaria, conjunctivitis, bronchospasm angioedema	Treated

The symptoms emerged after she consumed a fenugreek supplement to augment lactation while taking sertraline and breastfeeding.

Discussion

The utilization of herbal remedies, herbal medicinal products, and supplements has surged significantly over the past three decades. It is estimated that at least 80% of the world's population relies on these products for several aspects of healthcare, including preventive, therapeutic, or palliative care.^{85, 86} The use of herbal products is deeply entrenched in the traditional medicinal practices of many cultures for disease prevention and treatment.⁸⁷ Although the World Health Organization (WHO) acknowledges herbal products as a vital component of the healthcare system, growing apprehensions exist regarding their quality

and safety.⁸⁸ Herbal products are generally not subject to stringent regulation, as they are often classified as dietary supplements, thereby evading the rigorous scrutiny required for pharmaceutical drugs. Despite a widespread public perception that herbal remedies are inherently safe, numerous studies indicated that they could cause mild to severe, clinically significant adverse effects.^{28, 89}

Nowadays, the market for non-registered health-related products consists primarily of herbal remedies and food supplements. In this context, official or governmental systems often fail to comprehensively document their adverse effects, which are typically reported on a case-by-case basis. Research revealed that consumers might experience adverse events from herbal food supplements, particularly when they received insufficient information about these products.⁹⁰⁻⁹³

Fenugreek is commonly used as a food flavoring and is also consumed traditionally to prevent or treat certain diseases.^{94, 95} While numerous systematic reviews have examined its efficacy^{19, 96, 97} and toxicity,⁹⁸ no previous review has specifically focused on its adverse effects. This study is the first to systematically extract and summarize the adverse effects reported across various study types. This study illustrated the potential negative consequences associated with fenugreek consumption.

The most significant and well-documented adverse effects were allergic and hypersensitivity reactions, including angioedema, anaphylaxis, Stevens-Johnson syndrome, and toxic epidermal necrolysis. The clinical significance of these reactions is underscored by their documentation in dedicated case reports and case series. Other less common but notable adverse effects included serotonin syndrome, a maple syrup odor in infants, elevated INR level, dizziness, syncope, and other similar symptoms. However, the most frequently reported adverse effects were gastrointestinal problems, which were often mild and required no treatment. According to the included articles, all patients who experienced adverse effects from fenugreek recovered completely, with many cases resolving without intervention.

Several mechanisms could be suggested for the adverse effects associated with fenugreek. Gastrointestinal effects, such as indigestion, abdominal distention, and bloating, are likely attributable to its high fiber content.⁹⁹ This high fiber content may also interfere with the absorption of concurrently administered oral medications. Consequently, it is advisable to avoid combining fenugreek with oral drugs, particularly those with a narrow therapeutic index. The hypoglycemic activity of fenugreek, which occurs through the stimulation of insulin signaling pathways¹⁰⁰ and the inhibition of carbohydrate digestion and absorption by its soluble dietary fiber,¹⁰¹ poses a risk of hypoglycemia. Therefore, diabetic patients should use fenugreek with caution, and long-term use might necessitate an adjustment of their insulin dose. Moreover, fenugreek consumption was associated with reduced serum levels of potassium.⁷⁶ Evidence showed that when used concomitantly with diuretics or other hypokalemic agents, fenugreek might potentially contribute to hypokalemia.^{102, 103}

Dizziness associated with fenugreek use might result from a significant decrease in systolic blood pressure¹⁰¹ or from hypoglycemia-induced neurological symptoms due to its activity on insulin receptors.²² Moreover, concomitant use of fenugreek with antiarrhythmic drugs,

diuretics, laxatives, or mineralocorticoids might increase the risk of hypokalemia and potentiate cardiac glycoside toxicity. Fenugreek seed extracts exhibited both central nervous system (CNS) stimulant and depressant activities,^{77, 78} which could potentially trigger seizures in individuals with epilepsy.

The most frequently reported significant complications were allergic reactions to fenugreek.^{76-78, 81, 83, 84} As fenugreek belongs to the Fabaceae family, individuals with allergies to peanuts, peas, or coriander should use it with caution due to the potential for cross-reactivity. Fenugreek might also interact with serotonergic drugs, such as selective serotonin reuptake inhibitor (SSRI), increasing the risk of serotonin syndrome.⁸² A case report noted that fenugreek could elevate the INR, a critical consideration for patients taking warfarin.⁸⁰ A rare, benign, yet well-documented effect is a maple syrup odor in the urine or sweat of infants or mothers following maternal consumption.⁷⁹ Finally, the safety of long-term fenugreek use remains uncertain due to a lack of extensive studies.

This study had several limitations. As a scoping review, it did not incorporate critical steps of a systematic review, such as a formal quality assessment of the included studies, an evaluation of the risk of bias, or the use of independent reviewers for article selection and data extraction. Furthermore, the overall quality of the evidence was not assessed using a tool such as GRADE. The review was also limited by its exclusion of animal studies, experimental studies, case-control studies, cross-sectional studies, and cohort studies. Consequently, the authors recommend that future research consider these limitations.

It is important to note that a meta-analysis was not an objective of this study. In addition, while some of the included studies did not report any adverse events, this could not be taken as definitive proof of their absence. The authors suggested that a future systematic review and meta-analysis (including network meta-analysis) could provide more robust insights into the adverse effects of fenugreek and its supplements. Finally, further studies are strongly recommended to investigate the mechanisms underlying fenugreek's adverse effects and to assess the relationship between dosage, severity, and the incidence of these side effects.

Conclusion

Fenugreek and its supplements are commonly used worldwide as both a vegetable and an herbal remedy. Although widely perceived

as safe, several studies demonstrated that it could cause adverse effects, including allergic and hypersensitivity reactions, gastrointestinal complaints, headache, dizziness, hypoglycemia, elevated INR levels in plasma, and hypokalemia. However, it is crucial to note that no documented fatalities or cases requiring aggressive medical intervention were attributed to fenugreek use.

Acknowledgment

The authors extend their gratitude to the Vice-Chancellor for Research of Shiraz University of Medical Sciences (Shiraz, Iran) for supporting this study (Grant no: 95-01-01-11487). This manuscript is extracted from the doctoral thesis of Asma Homaei, which was completed in partial fulfillment of the requirements for the Medical Doctor degree at Shiraz University of Medical Sciences.

Authors' Contribution

E.AA: Study design, conception, data gathering, data analysis, data interpretation, drafting and reviewing critically; MM.P: Study design, data analysis, data interpretation, and reviewing critically; R.Z: Data analysis and drafting; M.MJ: Data gathering and drafting; Th.R: Study design and reviewing critically; M.P: Study design, conception, data gathering, data analysis, data interpretation and reviewing critically; All authors approved the final the version of the manuscript and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Declaration of AI

Artificial intelligence (QuillBot) was used solely for language editing. The authors are fully responsible for the content and integrity of the manuscript.

Conflict of Interest: None declared.

References

- 1 Jackson AM, Mullican L, Grizzle JH, Sullivan KL. Complementary and alternative medicine prevalence and patterns in the South-eastern United States. *Epidemiology Faculty and Staff Presentations*. 2016.
- 2 Harris PE, Cooper KL, Relton C, Thomas KJ. Prevalence of complementary and alternative medicine (CAM) use by the general population: a systematic review and update. *Int J Clin Pract*. 2012;66:924-39. doi: 10.1111/j.1742-1241.2012.02945.x. PubMed PMID: 22994327.
- 3 de Moraes Mello Boccolini P, Siqueira Boccolini C. Prevalence of complementary and alternative medicine (CAM) use in Brazil. *BMC Complement Med Ther*. 2020;20:51. doi: 10.1186/s12906-020-2842-8. PubMed PMID: 32054461; PubMed Central PMCID: PMC7076835.
- 4 Lee EL, Richards N, Harrison J, Barnes J. Prevalence of Use of Traditional, Complementary and Alternative Medicine by the General Population: A Systematic Review of National Studies Published from 2010 to 2019. *Drug Saf*. 2022;45:713-35. doi: 10.1007/s40264-022-01189-w. PubMed PMID: 35788539; PubMed Central PMCID: PMC9296440.
- 5 Parvizi MM, Forouhari S, Shahriarirad R, Shahriarirad S, Bradley RD, Roosta L. Prevalence and associated factors of complementary and integrative medicine use in patients afflicted with COVID-19. *BMC Complement Med Ther*. 2022;22:251. doi: 10.1186/s12906-022-03722-x. PubMed PMID: 36180868; PubMed Central PMCID: PMC9524323.
- 6 Bahall M. Complementary and alternative medicine usage among cardiac patients: a descriptive study. *BMC Complement Altern Med*. 2015;15:100. doi: 10.1186/s12906-015-0610-y. PubMed PMID: 25888160; PubMed Central PMCID: PMC4394556.
- 7 Posadzki P, Lee MS, Moon TW, Choi TY, Park TY, Ernst E. Prevalence of complementary and alternative medicine (CAM) use by menopausal women: a systematic review of surveys. *Maturitas*. 2013;75:34-43. doi: 10.1016/j.maturitas.2013.02.005. PubMed PMID: 23497959.
- 8 Slavin SL, Rogers RG, Komesu Y, Omotosho T, Hammil S, Lewis C, et al. Complementary and alternative medicine (CAM) use in women with pelvic floor disorders: a cohort study. *Int Urogynecol J*. 2010;21:431-7. doi: 10.1007/s00192-009-1058-8. PubMed PMID: 19967336; PubMed Central PMCID: PMC3729340.
- 9 Parvizi MM, Fatehi N, Jaladat AM, Gholampour Z, Shahriarirad R, Erfani A. Epidemiological factors in patients with dermatologic conditions referring to the clinic of traditional Persian Medicine: A cross-sectional study. *Int J Clin Pract*. 2021;75:e14788. doi: 10.1111/ijcp.14788. PubMed PMID: 34480839.
- 10 Hashempour MH, Mosavat SH, Heydari M, Shams M. Medicinal plants' use among

- patients with dyslipidemia: an Iranian cross-sectional survey. *J Complement Integr Med*. 2018;16. doi: 10.1515/jcim-2018-0101. PubMed PMID: 30391934.
- 11 Żuk-Golaszewska K, Wierzbowska J. Fenugreek: productivity, nutritional value and uses. *J Elementol*. 2017;22:1396-1411. doi: 10.5601/jelem.2017.22.1.1396.
 - 12 Aasim M, Baloch FS, Nadeem MA, Bakhsh A, Sameeullah M, Day S. Fenugreek (*Trigonella foenum-graecum* L.): an underutilized edible plant of modern world. In: Ozturk M, Hakeem KR, Ashraf M, Ahmad M, eds. *Global Perspectives on Underutilized Crops*. Cham: Springer; 2018. p. 381-8. doi: 10.1007/978-3-319-77776-4_12.
 - 13 Bahmani M, Shirzad H, Mirhosseini M, Mesripour A, Rafieian-Kopaei M. A Review on Ethnobotanical and Therapeutic Uses of Fenugreek (*Trigonella foenum-graecum* L.). *J Evid Based Complementary Altern Med*. 2016;21:53-62. doi: 10.1177/2156587215583405. PubMed PMID: 25922446.
 - 14 Singh P, Bajpai V, Gond V, Kumar A, Tadi-goppula N, Kumar B. Determination of Bioactive Compounds of Fenugreek (*Trigonella foenum-graecum*) Seeds Using LC-MS Techniques. *Methods Mol Biol*. 2020;2107:377-93. doi: 10.1007/978-1-0716-0235-5_21. PubMed PMID: 31893460.
 - 15 Malik A, Jamil U, Butt TT, Waquar S, Gan SH, Shafique H, et al. In silico and in vitro studies of lupeol and iso-orientin as potential antidiabetic agents in a rat model. *Drug Des Devel Ther*. 2019;13:1501-13. doi: 10.2147/DDDT.S176698. PubMed PMID: 31123393; PubMed Central PMCID: PMC6510393.
 - 16 Gupta RS, Grover AS, Kumar P, Goel A, Banik SP, Chakraborty S, et al. A randomized double blind placebo controlled trial to assess the safety and efficacy of a patented fenugreek (*Trigonella foenum-graecum*) seed extract in Type 2 diabetics. *Food Nutr Res*. 2024;68. doi: 10.29219/fnr.v68.10667. PubMed PMID: 38863744; PubMed Central PMCID: PMC11165257.
 - 17 Liu Y, Kakani R, Nair MG. Compounds in functional food fenugreek spice exhibit anti-inflammatory and antioxidant activities. *Food Chem*. 2012;131:1187-92. doi: 10.1016/j.foodchem.2011.09.102.
 - 18 Leela N, Shafeekh K. *Fenugreek, Chemistry of Spices*. Wallingford: CAB International; 2008. p. 242-51. doi: 10.1079/9781845934057.0242.
 - 19 El Nasri NA, El Tinay AH. Functional properties of fenugreek (*Trigonella foenum graecum*) protein concentrate. *Food Chem*. 2007;103:582-9. doi: 10.1016/j.foodchem.2006.09.003.
 - 20 Fatima T, Maqbool K, Hussain SZ. Potential health benefits of fenugreek. *J Med Plants Stud*. 2018;6:166-9.
 - 21 Akbari S, Abdurahman NH, Yunus RM, Alara OR, Abayomi OO. Extraction, characterization and antioxidant activity of fenugreek (*Trigonella foenum graecum*) seed oil. *Mater Sci Energy Technol*. 2019;2:349-55. doi: 10.1016/j.mset.2018.12.001.
 - 22 Haber SL, Keonavong J. Fenugreek use in patients with diabetes mellitus. *Am J Health Syst Pharm*. 2013;70:1196, 8, 200, 202-3. doi: 10.2146/ajhp120523. PubMed PMID: 23820455.
 - 23 Raju J, Patlolla JM, Swamy MV, Rao CV. Diosgenin, a steroid saponin of *Trigonella foenum graecum* (Fenugreek), inhibits azoxymethane-induced aberrant crypt foci formation in F344 rats and induces apoptosis in HT-29 human colon cancer cells. *Cancer Epidemiol Biomarkers Prev*. 2004;13:1392-8. PubMed PMID: 15298963.
 - 24 Sharma RD, Raghuram TC, Rao VD. Hypolipidaemic effect of fenugreek seeds. A clinical study. *Phytother Res*. 1991;5:145-7. doi: 10.1002/ptr.2650050313.
 - 25 Mansoori A, Hosseini S, Zilaei M, Hormoznejad R, Fathi M. Effect of fenugreek extract supplement on testosterone levels in male: A meta-analysis of clinical trials. *Phytother Res*. 2020;34:1550-5. doi: 10.1002/ptr.6627. PubMed PMID: 32048383.
 - 26 Navarro Del Hierro J, Casado-Hidalgo G, Reglero G, Martin D. The hydrolysis of saponin-rich extracts from fenugreek and quinoa improves their pancreatic lipase inhibitory activity and hypocholesterolemic effect. *Food Chem*. 2021;338:128113. doi: 10.1016/j.foodchem.2020.128113. PubMed PMID: 33092009.
 - 27 Ghasemi V, Kheirkhah M, Vahedi M. The Effect of Herbal Tea Containing Fenugreek Seed on the Signs of Breast Milk Sufficiency in Iranian Girl Infants. *Iran Red Crescent Med J*. 2015;17:e21848. doi: 10.5812/ircmj.21848. PubMed PMID: 26430522; PubMed Central PMCID: PMC4585338.
 - 28 Boullata JI, Nace AM. Safety issues with herbal medicine. *Pharmacotherapy*. 2000;20:257-69. doi: 10.1592/phco.20.4.257.34886. PubMed PMID: 10730682.
 - 29 Molavi Vardanjani H, Salehi Z, Alembizar F, Cramer H, Pasalar M. Prevalence and the Determinants of Traditional, Complementary, and Integrative Medicine Use

- Among Breastfeeding Mothers: A Cross-Sectional Study. *J Integr Complement Med*. 2022;28:67-76. doi: 10.1089/jicm.2021.0270. PubMed PMID: 35085020.
- 30 Abdel-Barry JA, Abdel-Hassan IA, Jawad AM, al-Hakim MH. Hypoglycaemic effect of aqueous extract of the leaves of *Trigonella foenum-graecum* in healthy volunteers. *East Mediterr Health J*. 2000;6:83-8. PubMed PMID: 11370345.
- 31 Rao A, Grant R. The effect of *Trigonella foenum-graecum* extract on prostate-specific antigen, and prostate function in otherwise healthy men with benign prostate hyperplasia. *Phytother Res*. 2020;34:634-9. doi: 10.1002/ptr.6554. PubMed PMID: 31828857.
- 32 Bumrungpert A, Somboonpanyakul P, Pavadhgul P, Thaninthanon S. Effects of Fenugreek, Ginger, and Turmeric Supplementation on Human Milk Volume and Nutrient Content in Breastfeeding Mothers: A Randomized Double-Blind Controlled Trial. *Breastfeed Med*. 2018;13:645-50. doi: 10.1089/bfm.2018.0159. PubMed PMID: 30411974.
- 33 Chevassus H, Molinier N, Costa F, Galtier F, Renard E, Petit P. A fenugreek seed extract selectively reduces spontaneous fat consumption in healthy volunteers. *Eur J Clin Pharmacol*. 2009;65:1175-8. doi: 10.1007/s00228-009-0733-5. PubMed PMID: 19809809.
- 34 Sharma RD, Raghuram TC, Rao NS. Effect of fenugreek seeds on blood glucose and serum lipids in type I diabetes. *Eur J Clin Nutr*. 1990;44:301-6. PubMed PMID: 2194788.
- 35 Gupta A, Gupta R, Lal B. Effect of *Trigonella foenum-graecum* (fenugreek) seeds on glycaemic control and insulin resistance in type 2 diabetes mellitus: a double blind placebo controlled study. *J Assoc Physicians India*. 2001;49:1057-61. PubMed PMID: 11868855.
- 36 Steels E, Rao A, Vitetta L. Physiological aspects of male libido enhanced by standardized *Trigonella foenum-graecum* extract and mineral formulation. *Phytother Res*. 2011;25:1294-300. doi: 10.1002/ptr.3360. PubMed PMID: 21312304.
- 37 Rao A, Steels E, Beccaria G, Inder WJ, Vitetta L. Influence of a Specialized *Trigonella foenum-graecum* Seed Extract (Libifem), on Testosterone, Estradiol and Sexual Function in Healthy Menstruating Women, a Randomised Placebo Controlled Study. *Phytother Res*. 2015;29:1123-30. doi: 10.1002/ptr.5355. PubMed PMID: 25914334.
- 38 Najdi RA, Hagraas MM, Kamel FO, Magadmi RM. A randomized controlled clinical trial evaluating the effect of *Trigonella foenum-graecum* (fenugreek) versus glibenclamide in patients with diabetes. *Afr Health Sci*. 2019;19:1594-601. doi: 10.4314/ahs.v19i1.34. PubMed PMID: 31148988; PubMed Central PMCID: PMC6531936.
- 39 Chevassus H, Gaillard JB, Farret A, Costa F, Gabillaud I, Mas E, et al. A fenugreek seed extract selectively reduces spontaneous fat intake in overweight subjects. *Eur J Clin Pharmacol*. 2010;66:449-55. doi: 10.1007/s00228-009-0770-0. PubMed PMID: 20020282.
- 40 Emtiazy M, Oveidzadeh L, Habibi M, Molaeipour L, Talei D, Jafari Z, et al. Investigating the effectiveness of the *Trigonella foenum-graecum* L. (fenugreek) seeds in mild asthma: a randomized controlled trial. *Allergy Asthma Clin Immunol*. 2018;14:19. doi: 10.1186/s13223-018-0238-9. PubMed PMID: 29743896; PubMed Central PMCID: PMC5930943.
- 41 Lu FR, Shen L, Qin Y, Gao L, Li H, Dai Y. Clinical observation on *trigonella foenum-graecum* L. total saponins in combination with sulfonylureas in the treatment of type 2 diabetes mellitus. *Chin J Integr Med*. 2008;14:56-60. doi: 10.1007/s11655-007-9005-3. PubMed PMID: 18219452.
- 42 Pandaran Sudheeran S, Jacob D, Natinga Mulakal J, Gopinathan Nair G, Maliakel A, Maliakel B, et al. Safety, Tolerance, and Enhanced Efficacy of a Bioavailable Formulation of Curcumin With Fenugreek Dietary Fiber on Occupational Stress: A Randomized, Double-Blind, Placebo-Controlled Pilot Study. *J Clin Psychopharmacol*. 2016;36:236-43. doi: 10.1097/JCP.0000000000000508. PubMed PMID: 27043120.
- 43 Mirgaloybayat S, Akbari Sene A, Jayervand F, Vazirian M, Mohazzab A, Kazerooni M. Comparison of the Effect of Fenugreek and Metformin on Clinical and Metabolic Status of Cases with Polycystic Ovary Syndrome: A Randomized Trial. *J Reprod Infertil*. 2024;25:120-32. doi: 10.18502/jri.v25i2.16040. PubMed PMID: 39157801; PubMed Central PMCID: PMC11327428.
- 44 Shamshad Begum S, Jayalakshmi HK, Vidyavathi HG, Gopakumar G, Abin I, Balu M, et al. A Novel Extract of Fenugreek Husk (FenuSMART) Alleviates Postmenopausal Symptoms and Helps to Establish the Hormonal Balance: A Randomized, Double-Blind, Placebo-Controlled Study. *Phytother Res*. 2016;30:1775-84. doi: 10.1002/ptr.5680. PubMed PMID: 27406028.
- 45 Swaroop A, Jaipuria AS, Gupta SK, Bagchi

- M, Kumar P, Preuss HG, et al. Efficacy of a Novel Fenugreek Seed Extract (*Trigonella foenum-graecum*, Furocyst) in Polycystic Ovary Syndrome (PCOS). *Int J Med Sci*. 2015;12:825-31. doi: 10.7150/ijms.13024. PubMed PMID: 26516311; PubMed Central PMCID: PMC4615243.
- 46 Nathan J, Panjwani S, Mohan V, Joshi V, Thakurdesai PA. Efficacy and safety of standardized extract of *Trigonella foenum-graecum* L seeds as an adjuvant to L-Dopa in the management of patients with Parkinson's disease. *Phytother Res*. 2014;28:172-8. doi: 10.1002/ptr.4969. PubMed PMID: 23512705.
 - 47 Florentin M, Liberopoulos E, Elisaf MS, Tsimihodimos V. No effect of fenugreek, bergamot and olive leaf extract on glucose homeostasis in patients with prediabetes: a randomized double-blind placebo-controlled study. *Arch Med Sci Atheroscler Dis*. 2019;4:e162-e6. doi: 10.5114/amsad.2019.86756. PubMed PMID: 31448348; PubMed Central PMCID: PMC6704765.
 - 48 Verma N, Usman K, Patel N, Jain A, Dhakre S, Swaroop A, et al. A multicenter clinical study to determine the efficacy of a novel fenugreek seed (*Trigonella foenum-graecum*) extract (Fenfurol) in patients with type 2 diabetes. *Food Nutr Res*. 2016;60:32382. doi: 10.3402/fnr.v60.32382. PubMed PMID: 27733237; PubMed Central PMCID: PMC5061863.
 - 49 Geberemeskel GA, Debebe YG, Nguse NA. Antidiabetic Effect of Fenugreek Seed Powder Solution (*Trigonella foenum-graecum* L.) on Hyperlipidemia in Diabetic Patients. *J Diabetes Res*. 2019;2019:8507453. doi: 10.1155/2019/8507453. PubMed PMID: 31583253; PubMed Central PMCID: PMC6748210.
 - 50 Madar Z, Abel R, Samish S, Arad J. Glucose-lowering effect of fenugreek in non-insulin dependent diabetics. *Eur J Clin Nutr*. 1988;42:51-4. PubMed PMID: 3286242.
 - 51 Younesy S, Amiraliakbari S, Esmaeili S, Alavimajd H, Nouraei S. Effects of fenugreek seed on the severity and systemic symptoms of dysmenorrhea. *J Reprod Infertil*. 2014;15:41-8. PubMed PMID: 24695380; PubMed Central PMCID: PMC3955423.
 - 52 Losso JN, Holliday DL, Finley JW, Martin RJ, Rood JC, Yu Y, et al. Fenugreek bread: a treatment for diabetes mellitus. *J Med Food*. 2009;12:1046-9. doi: 10.1089/jmf.2008.0199. PubMed PMID: 19857068.
 - 53 Inanmdar W, Sultana A, Mubeen U, Rahman K. Clinical efficacy of *Trigonella foenum graecum* (Fenugreek) and dry cupping therapy on intensity of pain in patients with primary dysmenorrhea. *Chin J Integr Med*. 2016. doi: 10.1007/s11655-016-2259-x. PubMed PMID: 27225291.
 - 54 Park HJ, Lee KS, Lee EK, Park NC. Efficacy and Safety of a Mixed Extract of *Trigonella foenum-graecum* Seed and *Lespedeza cuneata* in the Treatment of Testosterone Deficiency Syndrome: A Randomized, Double-Blind, Placebo-Controlled Clinical Trial. *World J Mens Health*. 2018;36:230-8. doi: 10.5534/wjmh.170004. PubMed PMID: 29623697; PubMed Central PMCID: PMC6119847.
 - 55 Hassanzadeh Bashtian M, Emami SA, Mousavifar N, Esmaily HA, Mahmoudi M, Mohammad Poor AH. Evaluation of Fenugreek (*Trigonella foenum-graecum* L.), Effects Seeds Extract on Insulin Resistance in Women with Polycystic Ovarian Syndrome. *Iran J Pharm Res*. 2013;12:475-81. PubMed PMID: 24250624; PubMed Central PMCID: PMC3813238.
 - 56 Rao A, Steels E, Inder WJ, Abraham S, Vitetta L. Testofen, a specialised *Trigonella foenum-graecum* seed extract reduces age-related symptoms of androgen decrease, increases testosterone levels and improves sexual function in healthy aging males in a double-blind randomised clinical study. *Aging Male*. 2016;19:134-42. doi: 10.3109/13685538.2015.1135323. PubMed PMID: 26791805.
 - 57 Maheshwari A, Verma N, Swaroop A, Bagchi M, Preuss HG, Tiwari K, et al. Efficacy of Furosap(TM), a novel *Trigonella foenum-graecum* seed extract, in Enhancing Testosterone Level and Improving Sperm Profile in Male Volunteers. *Int J Med Sci*. 2017;14:58-66. doi: 10.7150/ijms.17256. PubMed PMID: 28138310; PubMed Central PMCID: PMC5278660.
 - 58 Hausenblas HA, Conway KL, Coyle KRM, Barton E, Smith LD, Esposito M, et al. Efficacy of fenugreek seed extract on men's psychological and physical health: a randomized placebo-controlled double-blind clinical trial. *J Complement Integr Med*. 2020;18:445-8. doi: 10.1515/jcim-2019-0101. PubMed PMID: 32441668.
 - 59 Zarghi A, Haddadi M, Tabarraie Y, Movahedzadeh D, Ghobadi F, Sarpooshi HR. Effect of fenugreek (*Trigonella foenum-graecum* L.) seeds powder on gastrointestinal bleeding in mechanically ventilated patients: A double-blind, randomized controlled clinical trial. *J Tradit Chin Med Sci*. 2021;8:150-4. doi: 10.1016/j.jtcms.2021.05.003.

- 60 Foroumandi E, Javan R, Moayed L, Fahimi H, Kheirabadi F, Neamatshahi M, et al. The effects of fenugreek seed extract supplementation in patients with Alzheimer's disease: A randomized, double-blind, placebo-controlled trial. *Phytother Res.* 2023;37:285-94. doi: 10.1002/ptr.7612. PubMed PMID: 36199177.
- 61 Lee-Odegard S, Gundersen TE, Drevon CA. Effect of a plant extract of fenugreek (*Trigonella foenum-graecum*) on testosterone in blood plasma and saliva in a double blind randomized controlled intervention study. *PLoS One.* 2024;19:e0310170. doi: 10.1371/journal.pone.0310170. PubMed PMID: 39288153; PubMed Central PMCID: PMC11407615.
- 62 Hota D, Padhy BM, Maiti R, Bisoi D, Sahoo JP, Patro BK, et al. A Placebo-Controlled, Double-Blind Clinical Investigation to Evaluate the Efficacy of a Patented *Trigonella foenum-graecum* Seed Extract "Fenfuro(R)" in Type 2 Diabetics. *J Am Nutr Assoc.* 2024;43:147-56. doi: 10.1080/27697061.2023.2233008. PubMed PMID: 37459747.
- 63 Singh A, Gainer S, Banerjee P, Goel A, Kumar P, Mondal B, et al. Efficacy of a Proprietary Fenugreek Seed Extract (*Trigonella foenum-graecum*, Furocyst(R)) in Women with Polycystic Ovary Syndrome (PCOS): a Randomized, Double-Blind, Placebo-Controlled Study. *J Am Nutr Assoc.* 2023;42:651-9. doi: 10.1080/27697061.2022.2126410. PubMed PMID: 36219198.
- 64 Robert SD, Ismail AA, Rosli WI. Reduction of postprandial blood glucose in healthy subjects by buns and flatbreads incorporated with fenugreek seed powder. *Eur J Nutr.* 2016;55:2275-80. doi: 10.1007/s00394-015-1037-4. PubMed PMID: 26358163.
- 65 Sundaram G, Ramakrishnan T, Parthasarathy H, Raja M, Raj S. Fenugreek, diabetes, and periodontal disease: A cross-link of sorts! *J Indian Soc Periodontol.* 2018;22:122-6. doi: 10.4103/jisp.jisp_322_17. PubMed PMID: 29769766; PubMed Central PMCID: PMC5939019.
- 66 Ansari M, Sadeghi P, Mahdavi H, Fattahi-Dolatabadi M, Mohamadi N, Asadi A, et al. Fenugreek dermal patch, a new natural topical antinociceptive medication for relieving the postherniotomy pain, a double-blind placebo controlled trial. *J Complement Integr Med.* 2019;16. doi: 10.1515/jcim-2018-0082. PubMed PMID: 30721143.
- 67 Bordia A, Verma SK, Srivastava KC. Effect of ginger (*Zingiber officinale* Rosc.) and fenugreek (*Trigonella foenum-graecum* L.) on blood lipids, blood sugar and platelet aggregation in patients with coronary artery disease. *Prostaglandins Leukot Essent Fatty Acids.* 1997;56:379-84. doi: 10.1016/s0952-3278(97)90587-1. PubMed PMID: 9175175.
- 68 Kassaian N, Azadbakht L, Forghani B, Amini M. Effect of fenugreek seeds on blood glucose and lipid profiles in type 2 diabetic patients. *Int J Vitam Nutr Res.* 2009;79:34-9. doi: 10.1024/0300-9831.79.1.34. PubMed PMID: 19839001.
- 69 Ranade M, Mudgalkar N. A simple dietary addition of fenugreek seed leads to the reduction in blood glucose levels: A parallel group, randomized single-blind trial. *Ayu.* 2017;38:24-7. doi: 10.4103/ayu.AYU_209_15. PubMed PMID: 29861588; PubMed Central PMCID: PMC5954247.
- 70 Kiss R, Szabo K, Gesztelyi R, Somodi S, Kovacs P, Szabo Z, et al. Insulin-Sensitizer Effects of Fenugreek Seeds in Parallel with Changes in Plasma MCH Levels in Healthy Volunteers. *Int J Mol Sci.* 2018;19. doi: 10.3390/ijms19030771. PubMed PMID: 29518003; PubMed Central PMCID: PMC5877632.
- 71 Mathern JR, Raatz SK, Thomas W, Slavin JL. Effect of fenugreek fiber on satiety, blood glucose and insulin response and energy intake in obese subjects. *Phytother Res.* 2009;23:1543-8. doi: 10.1002/ptr.2795. PubMed PMID: 19353539.
- 72 Steels E, Steele ML, Harold M, Coulson S. Efficacy of a Proprietary *Trigonella foenum-graecum* L. De-Husked Seed Extract in Reducing Menopausal Symptoms in Otherwise Healthy Women: A Double-Blind, Randomized, Placebo-Controlled Study. *Phytother Res.* 2017;31:1316-22. doi: 10.1002/ptr.5856. PubMed PMID: 28707431.
- 73 Palacios S, Soler E, Ramirez M, Lilue M, Khorsandi D, Losa F. Effect of a multi-ingredient based food supplement on sexual function in women with low sexual desire. *BMC Womens Health.* 2019;19:58. doi: 10.1186/s12905-019-0755-9. PubMed PMID: 31039769; PubMed Central PMCID: PMC6492381.
- 74 Mehrzadi S, Mirzaei R, Heydari M, Sasani M, Yaqoobvand B, Huseini HF. Efficacy and Safety of a Traditional Herbal Combination in Patients with Type II Diabetes Mellitus: A Randomized Controlled Trial. *J Diet Suppl.* 2021;18:31-43. doi: 10.1080/19390211.2020.1727076. PubMed PMID: 32081056.
- 75 Varghese N, Ramesh A, Potdar R. Clinical Evaluation of Fenugreek Toothpaste and

- Regular Toothpaste in Control of Gingivitis—A Comparative Study. *Int J Sci Res Publ.* 2020;13:133-7.
- 76 Patil SP, Niphadkar PV, Bapat MM. Allergy to fenugreek (*Trigonella foenum graecum*). *Ann Allergy Asthma Immunol.* 1997;78:297-300. doi: 10.1016/S1081-1206(10)63185-7. PubMed PMID: 9087156.
 - 77 Bentele-Jaberg N, Guenova E, Mehra T, Nageli M, Chang YT, Cozzio A, et al. The Phytotherapeutic Fenugreek as Trigger of Toxic Epidermal Necrolysis. *Dermatology.* 2015;231:99-102. doi: 10.1159/000433423. PubMed PMID: 26138328.
 - 78 Joseph NI, Slavin E, Peppers BP, Hostoffer RW, Jr. Fenugreek Anaphylaxis in a Pediatric Patient. *Allergy Rhinol (Providence).* 2018;9:2152656718764134. doi: 10.1177/2152656718764134. PubMed PMID: 29977649; PubMed Central PMCID: PMC6028160.
 - 79 Korman SH, Cohen E, Preminger A. Pseudo-maple syrup urine disease due to maternal prenatal ingestion of fenugreek. *J Paediatr Child Health.* 2001;37:403-4. doi: 10.1046/j.1440-1754.2001.00617.x. PubMed PMID: 11532065.
 - 80 Lambert JP, Cormier J. Potential interaction between warfarin and boldo-fenugreek. *Pharmacotherapy.* 2001;21:509-12. doi: 10.1592/phco.21.5.509.34492. PubMed PMID: 11310527.
 - 81 Aurich S, Spiric J, Engin A, Simon JC, Mahler V, Treudler R. Report of a Case of IgE-Mediated Anaphylaxis to Fenugreek. *J Investig Allergol Clin Immunol.* 2019;29:56-8. doi: 10.18176/jiaci.0328. PubMed PMID: 30785103.
 - 82 Doolabh K, Finnegan D, Pehlivan N, Farrand S. Oral fenugreek seed consumption and serotonin syndrome. *Aust N Z J Psychiatry.* 2019;53:1225. doi: 10.1177/0004867419866912. PubMed PMID: 31379190.
 - 83 Ohnuma N, Yamaguchi E, Kawakami Y. Anaphylaxis to curry powder. *Allergy.* 1998;53:452-4. doi: 10.1111/j.1398-9995.1998.tb03924.x. PubMed PMID: 9574894.
 - 84 Ebo DG, Bridts CH, Mertens MH, Stevens WJ. Coriander anaphylaxis in a spice grinder with undetected occupational allergy. *Acta Clin Belg.* 2006;61:152-6. doi: 10.1179/acb.2006.025. PubMed PMID: 16881566.
 - 85 Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol.* 2014;4:177. doi: 10.3389/fphar.2013.00177. PubMed PMID: 24454289; PubMed Central PMCID: PMC3887317.
 - 86 Suna S, Tamer CE, Özcan-Sinir G. Trends and possibilities of the usage of medicinal herbal extracts in beverage production. In: Grumezescu AM, Holban AM, eds. *Natural Beverages.* London: Academic Press; 2019. p. 361-98.
 - 87 Karunamoorthi K, Jegajeevanram K, Vijayalakshmi J, Mengistie E. Traditional medicinal plants: a source of phytotherapeutic modality in resource-constrained health care settings. *J Evid Based Complement Altern Med.* 2013;18:67-74. doi: 10.1177/2156587212460241.
 - 88 Onakpoya IJ, Heneghan CJ, Aronson JK. Post-marketing withdrawal of 462 medicinal products because of adverse drug reactions: a systematic review of the world literature. *BMC Med.* 2016;14:10. doi: 10.1186/s12916-016-0553-2. PubMed PMID: 26843061; PubMed Central PMCID: PMC4740994.
 - 89 Zhang J, Onakpoya IJ, Posadzki P, Eddouks M. The safety of herbal medicine: from prejudice to evidence. *Evid Based Complement Alternat Med.* 2015;2015:316706. doi: 10.1155/2015/316706. PubMed PMID: 25838831; PubMed Central PMCID: PMC4370194.
 - 90 Timbo BB, Chirtel SJ, Ihrie J, Oladipo T, Velez-Suarez L, Brewer V, et al. Dietary Supplement Adverse Event Report Data From the FDA Center for Food Safety and Applied Nutrition Adverse Event Reporting System (CAERS), 2004-2013. *Ann Pharmacother.* 2018;52:431-8. doi: 10.1177/1060028017744316. PubMed PMID: 29171279.
 - 91 Lenssen KGM, Bast A, de Boer A. How does scientific information reach the consumer? A case study among students into providing verbal information on dietary supplements at point of purchase. *Int J Food Sci Nutr.* 2021;72:402-17. doi: 10.1080/09637486.2020.1817344. PubMed PMID: 32907414.
 - 92 Chiba T, Sato Y, Nakanishi T, Yokotani K, Suzuki S, Umegaki K. Inappropriate usage of dietary supplements in patients by miscommunication with physicians in Japan. *Nutrients.* 2014;6:5392-404. doi: 10.3390/nu6125392. PubMed PMID: 25431879; PubMed Central PMCID: PMC4276974.
 - 93 de Boer A, Geboers L, van de Koppel S, van Hunsel F. Governance of nutravigilance in the Netherlands: Reporting adverse events of non-registered products. *Health Policy.* 2022;126:731-7. doi: 10.1016/j.

- healthpol.2022.05.011. PubMed PMID: 35643570.
- 94 Khorshidian N, Yousefi Asli M, Arab M, Adeli Mirzaie A, Mortazavian AM. Fenugreek: potential applications as a functional food and nutraceutical. *Nutraceutical Nutr Food Sci Res.* 2016;3:5-16. doi: 10.18869/acad-pub.nfsr.3.1.5.
- 95 Jhahhria A, Kumar K. Fenugreek with its medicinal applications. *Int J Pharm Sci Rev Res.* 2016;41:194-201.
- 96 Kandhare AD, Thakurdesai PA, Wangikar P, Bodhankar SL. A systematic literature review of fenugreek seed toxicity by using ToxRTool: evidence from preclinical and clinical studies. *Heliyon.* 2019;5:e01536. doi: 10.1016/j.heliyon.2019.e01536. PubMed PMID: 31049444; PubMed Central PMCID: PMC6482331.
- 97 Vijayakumar MV, Singh S, Chhipa RR, Bhat MK. The hypoglycaemic activity of fenugreek seed extract is mediated through the stimulation of an insulin signalling pathway. *Br J Pharmacol.* 2005;146:41-8. doi: 10.1038/sj.bjp.0706312. PubMed PMID: 15980869; PubMed Central PMCID: PMC1576255.
- 98 Hannan JM, Ali L, Rokeya B, Khaleque J, Akhter M, Flatt PR, et al. Soluble dietary fibre fraction of *Trigonella foenum-graecum* (fenugreek) seed improves glucose homeostasis in animal models of type 1 and type 2 diabetes by delaying carbohydrate digestion and absorption, and enhancing insulin action. *Br J Nutr.* 2007;97:514-21. doi: 10.1017/S0007114507657869. PubMed PMID: 17313713.
- 99 Hadi A, Arab A, Hajianfar H, Talaei B, Miraghajani M, Babajafari S, et al. The effect of fenugreek seed supplementation on serum irisin levels, blood pressure, and liver and kidney function in patients with type 2 diabetes mellitus: A parallel randomized clinical trial. *Complement Ther Med.* 2020;49:102315. doi: 10.1016/j.ctim.2020.102315. PubMed PMID: 32147060.
- 100 Sauvaire Y, Ribes G, Baccou JC, Loubatieeres-Mariani MM. Implication of steroid saponins and sapogenins in the hypocholesterolemic effect of fenugreek. *Lipids.* 1991;26:191-7. doi: 10.1007/BF02543970. PubMed PMID: 2046485.
- 101 Natarajan B, Muralidharan A, Satish R, Dhananjayan R. Neuropharmacological Activity of *Trigonella foenum graecum* Linn. Seeds. *J Nat Remedies.* 2007;7:160-5.
- 102 Singletary KW. Fenugreek: overview of potential health benefits. *Nutr Today.* 2017;52:93-111. doi: 10.1097/NT.0000000000000209.
- 103 Smith M. Therapeutic applications of fenugreek. *Altern Med Rev.* 2003;8:20-7.