



From Gene Expression to Candidate Biomarkers in Ulcerative Colitis: Aquaporin-9 in Neutrophil Extracellular Traps Formation Pathways

Hawra Zia Hossein Al-Jamali¹, MSc; 
Farshad Sheikhesmaili², MD, PhD;
Mohammad Moradzad¹, MSc; Amjad
Ahmadi³, PhD; Ali Jalili⁴, PhD; Bahram
Nikkhoo^{4,5}, MD, PhD; Mohammad Reza
Rahmani¹, PhD; Shohreh Fakhari⁵, PhD 

¹Department of Immunology, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran;

²Liver and Digestive Research Center, Research Institute for Health Development, Kurdistan University of Medical Sciences, Sanandaj, Iran;

³Health Metrics and Evaluation Research Center, Research Institute for Health Development, Kurdistan University of Medical Sciences, Sanandaj, Iran;

⁴Cancer and Immunology Research Center, Research Institute for Health Development, Kurdistan University of Medical Sciences, Sanandaj, Iran;

⁵Cellular and Molecular Research Center, Research Institute for Health Development, Kurdistan University of Medical Sciences, Sanandaj, Iran

Correspondence:

Shohreh Fakhari, PhD;
Kurdistan University of Medical Sciences,
Pasdaran Blvd., Postal code: 66177-13446,
Sanandaj, Iran

Tel: +98 87 33664645

Email: sh.fakhari@muk.ac.ir

Received: 06 October 2025

Revised: 12 December 2025

Accepted: 15 January 2026

What's Known

- Ulcerative colitis (UC) involves chronic intestinal inflammation, where neutrophil extracellular traps (NETs) contribute to disease progression by disrupting the mucosal barrier. Excessive NET formation correlates with increased inflammation and disease severity. Despite their critical role, the detailed mechanisms driving NET formation and reliable tissue biomarkers in UC remain unclear.

What's New

- This study identifies *AQP9* as a highly upregulated gene in UC, validated in both new and treatment-resistant patients. *AQP9* shows strong diagnostic accuracy, distinguishing UC patients from healthy controls, and, along with *FPR1* and *FPR2*, may serve as key biomarkers and therapeutic targets in UC management.

Abstract

Background: Ulcerative colitis (UC) is a chronic inflammatory bowel disease (IBD). Neutrophil extracellular traps (NETs) have been recognized as contributing to UC progression. This study aimed to identify key genes driving NET formation in UC and evaluate their potential as biomarkers.

Methods: Gene expression profiles from colon biopsies of UC patients and healthy controls were obtained from the GSE224758 dataset. Differentially expressed genes (DEGs) were identified using the GEO2R tool. Hub genes involved in NET formation, including *FCGR3B*, *AQP9*, *FPR1*, *FPR2*, and *NCF2*, were validated through reverse transcription-quantitative polymerase chain reaction (RT-qPCR) using colon biopsy samples from healthy controls (n=20), newly diagnosed UC (n=20), and treatment-resistant UC patients (n=20) collected between November 8, 2022, and February 2, 2025, at Tohid Hospital, Kurdistan University of Medical Sciences, Sanandaj, Iran. Data normality was assessed using the Shapiro–Wilk test; Kruskal–Wallis test and Dunn's *post hoc* test were applied for group comparisons. ROC analysis was applied to determine the potential of genes for the differentiation of healthy people from patients.

Results: RT-qPCR validation confirmed significantly increased expression of *AQP9* (P < 0.0001), *FPR1*, and *FPR2* (P=0.03 and P=0.02) genes in patients with UC and those with refractory UC compared to healthy controls. Among these, *AQP9* exhibited the most significant differential expression, demonstrating high sensitivity and specificity in distinguishing newly diagnosed and treatment-resistant UC patients from healthy individuals (sensitivity 83.49% and 80.98%, and specificity 67.77% and 64.24%, respectively), indicating its potential as a diagnostic biomarker.

Conclusion: This study identifies NET-associated genes, particularly *AQP9*, *FPR1*, and *FPR2*, as candidate tissue biomarkers for UC, with *AQP9* exhibiting the strongest ability to distinguish patients from healthy controls. These findings support the utility of NET-related genes as tissue biomarkers for UC, while emphasizing that any therapeutic targeting of these molecules lies beyond the scope of the present work and will require additional functional and *in vivo* validation.

Please cite this article as: Al-Jamali HZH, Sheikhesmaili F, Moradzad M, Ahmadi A, Jalili A, Nikkhoo B, Rahmani MR, Fakhari S. From Gene Expression to Candidate Biomarkers in Ulcerative Colitis: Aquaporin-9 in Neutrophil Extracellular Traps Formation Pathways. Iran J Med Sci. 2026;51(6):426-436. doi: 10.30476/ijms.2026.109059.4425.

Keywords • Colitis, ulcerative • Neutrophil extracellular traps • *Aquaporin 9* • Gene expression profiling

Introduction

Ulcerative colitis (UC) is a chronic inflammatory bowel disease (IBD) characterized by recurring inflammation of the colon with periods of flare-ups and remission. It commonly affects individuals aged 30 to 40, with a rising global incidence and prevalence, including about 0.3% of the European population diagnosed with UC.^{1, 2} Symptoms observed include diarrhea, presence of blood in stool, decreased appetite, abdominal discomfort, nausea, vomiting, and weight loss.³ Disease severity varies from localized rectal inflammation to extensive colitis, and diagnosis involves clinical evaluation, laboratory tests, imaging, and colonoscopy. Treatment progresses through stages, including 5-aminosalicylic acid, glucocorticoids, azathioprine, and anti-Tumor Necrosis Factor (TNF) therapy. Refractory patients are defined by failure to respond to all therapeutic stages.⁴

The exact cause of inflammation in UC is unknown. Since the immune system mistakenly attacks the intestinal lining, causing inflammation, UC is considered an autoimmune disease. This inappropriate immune response may be triggered by intestinal bacteria or other environmental factors, such as diet, stress, and infections. Dysbiosis, or an imbalance in gut microbiota, might also contribute to colonic inflammation.^{5, 6} The pathogenesis of UC is complex and involves genetic, environmental, immune dysregulation, and microbiome alterations, leading to the recruitment of various inflammatory cells, including neutrophils, dendritic cells, T cells, and B cells, to the intestine.³

Neutrophils play a critical role in UC by contributing to tissue damage through the release of reactive oxygen species, proteases, and pro-inflammatory cytokines.⁷⁻⁹ An important neutrophil defense mechanism is NETosis, where neutrophils release neutrophil extracellular traps (NETs), web-like structures composed of DNA and antimicrobial proteins that trap pathogens.¹⁰ While NETs help contain infections, excessive or dysregulated NET formation promotes inflammation and tissue injury.¹¹⁻¹³ Increased NET components have been found in inflamed mucosa, stool, and blood of UC patients, correlating with disease activity.¹¹ Previous studies have conducted transcriptomic and bioinformatic profiling of NET-associated genes in UC, identifying candidates such as *CXCR1* and *CXCR2* as potential diagnostic or prognostic biomarkers. These studies used methods including Mendelian randomization, machine

learning, and co-expression network analyses to implicate NET-related genes in UC pathogenesis and response prediction.¹⁴⁻¹⁶ However, the molecular mechanisms driving NET formation in UC remain poorly understood.

In this study, we performed bioinformatic analysis of gene expression data from colonic biopsies of UC patients and healthy controls and identified DEGs involved in NET formation, including Fc Gamma Receptor IIIb (*FCGR3B*), Aquaporin-9 (*AQP9*), Formyl Peptide Receptor 1 (*FPR1*), Formyl Peptide Receptor 2 (*FPR2*), and Neutrophil cytosol factor 2 (*NCF2*). These five genes were specifically selected from the largest gene module identified through PPI network analysis of 429 upregulated DEGs from the GSE224758 dataset. Pathway enrichment analysis using DAVID and ShinyGO revealed significant enrichment of the NET formation pathway within this module, with these five genes representing the complete set annotated to NET formation by KEGG pathway analysis. *FCGR3B* is a receptor expressed on neutrophils that induces NET formation through cross-linking with specific monoclonal antibodies. *AQP9* regulates neutrophil migration and hyperactivation, promoting NET formation, with its distinct expression pattern highlighting biomarker potential. *FPR1* and *FPR2*, G protein-coupled receptors, activate neutrophils and NETosis by recognizing formylated peptides from bacteria and damage signals, with roles demonstrated in multiple inflammatory diseases. *NCF2*, an NADPH oxidase subunit, generates ROS essential for NETosis and has been associated with NET formation in inflammatory conditions.¹⁷⁻²⁵

By validating the expression of these key genes in new and treatment-resistant UC patients, we aim to elucidate their role in UC pathogenesis and assess their potential as tissue biomarkers for improved diagnosis and therapeutic targeting. Our findings may provide novel insights into the contribution of NETs in UC and suggest new avenues for managing difficult-to-treat cases.

Materials and Methods

RNA Sequencing (RNA-seq) Data Resources and Identification of DEGs

This study employed a two-pronged approach, combining a bioinformatic analysis of publicly available RNA-seq data with subsequent experimental validation using patient-derived samples in a cross-sectional study design. RNA-seq data from the GSE224758 dataset, gene expression profiling of colon biopsies

from UC patients and healthy volunteers, was obtained from the Gene Expression Omnibus (GEO) database (<https://www.ncbi.nlm.nih.gov/geo/>) to identify DEGs. The GEO2R online tool (<http://www.ncbi.nlm.nih.gov/geo/geo2r/>) (NCBI, USA) was used to identify DEGs between UC and healthy control groups. This analysis employed the limma package for differential expression analysis, incorporating normalization methods such as quantile normalization to ensure comparability across samples by adjusting for technical variations. When addressing multiple testing correction, the Bonferroni correction was applied to control for false positives ([supplementary file S1](#)). DEGs were filtered using adjusted P value <0.01 and Log₂ fold change threshold=1, yielding 492 DEGs (429 upregulated, 63 downregulated) in [supplementary file S2](#).

Protein-Protein Interaction Network and Clustering

For the DEGs that are upregulated in the UC group, a co-expression Protein-Protein Interaction (PPI) network was constructed using the STRING database (version 12.0; <https://string-db.org>, Switzerland) with a medium confidence score (0.4) ([supplementary file S3](#)). The network was exported and visualized using Cytoscape software (version 3.10.1; Cytoscape Consortium, USA) ([supplementary file S4](#)). Clustering and module identification were performed using the Clusterviz plugin with the Fast Agglomerate Algorithm based on Edge Clustering Coefficients (FAG-EC algorithm) and a complex size threshold ≥3, identifying 14 modules. The largest module contained 31 nodes and 83 edges.

Gene Ontology and Pathway Enrichment Analyses

Gene Ontology and KEGG pathway enrichment analysis were conducted for the genes in the largest module using the DAVID database (<https://davidbioinformatics.nih.gov>) (NCBI/NIAID, USA). Results for KEGG pathways were downloaded and visualized using ShinyGO 0.81 (version 0.81; South Dakota State University, USA). The NET formation pathway was significantly enriched with *FCGR3B*, *AQP9*, *FPR1*, *FPR2*, and *NCF2* genes.

Patient Characteristics and Sample Collection

The study population and sample collection methodology here were based on our previous study.²⁶ In brief, colon biopsy samples were collected between November 8, 2022, and February 2, 2025, at Tohid Hospital (Kurdistan University of Medical Sciences, Sanandaj, Iran). Written informed consent was secured from every participant involved in the study. Three groups

were included: healthy controls (n=20), newly diagnosed UC patients (n=20), and treatment-resistant UC patients (n=20). Healthy controls had suspected UC ruled out by colonoscopy and pathology. UC diagnosis followed clinical, laboratory, endoscopic (Mayo score ≥6), and pathological criteria by gastroenterology specialists. Treatment-resistant UC was defined as failure to respond to 5-aminosalicylic acid (5-ASA), glucocorticoids, azathioprine, and anti-TNF therapy. Exclusion criteria included diabetes, rheumatoid arthritis, irritable bowel syndrome (IBS), and immunodeficiency. Colon biopsy samples were immediately placed in RNeasy lysis buffer (Qiagen, USA) and stored at -70 °C. The study protocol was thoroughly reviewed and approved by the Ethics Committee at Kurdistan University of Medical Sciences, under approval number IR.MUK.REC.1403.285.

RNA Extraction and Synthesis of cDNA

RNA extraction was performed using the FavorPrep™ Total RNA Mini Kit (Cat. No. FABRK 000-Mini, Favorgen Biotech, Taiwan) following the manufacturer's instructions. RNA quality was assessed (A260/280 1.8-2.0), and the purified RNA was stored at -70 °C until further analysis. Complementary DNA (cDNA) was synthesized from 1 µg RNA using the YT4500 cDNA Synthesis Kit (Yekta Tajhiz Azma, Iran) according to the standard protocol. The synthesized cDNA was diluted 1:5 and stored at -20 °C.

Reverse Transcription-Quantitative Polymerase Chain Reaction (RT-qPCR)

For the RT-qPCR assay, primers were designed using Gene Runner software (Hastings Software, USA), as listed in table 1. Each 20 µL reaction mixture contained 10 µL of YTA SYBR Green qPCR Mastermix 2X (Cat. No. YT2551; Yekta Tajhiz Azma, Iran), 1 µL each of forward and reverse primers (10 µM), 2 µL of cDNA template, and 6 µL of nuclease-free water. The thermal cycling conditions were carried out according to the manufacturer's protocol after preparing the reaction mixture.

To confirm primer specificity and optimize reaction conditions, a temperature gradient PCR was conducted for each primer pair targeting *FCGR3B*, *AQP9*, *FPR1*, *FPR2*, *NCF2*, and the reference gene β-ACTB. PCR products were analyzed on 2% agarose gels to verify the presence of single amplicons of expected sizes (table 1 and [supplementary figure 1S](#)). Standard curves generated from 5-point, 10-fold serial cDNA dilutions confirmed amplification efficiencies (E) of 90-110% for all primers.

Table 1: Specific primers to assess the genes involved in NETs formation and internal control

| Primer | Accession codes | Forward Sequence (5' to 3') | Reverse Sequence (5' to 3') | Amplicon Size (bp) |
|---------------|---|-----------------------------|-----------------------------|--------------------|
| β -ACTB | NM_001101 | AGATCATTGCTCCTCCTGAG | AGTCATAGTCCGCCTAGAAG | 159 |
| FCGR3B | NM_001244753 NM_000570 NM_001271035 NM_001271036 NM_001271037 | TCCTGGAGCCTCAATGGTAC | TGGCAGCGTCAATGAAGTAG | 153 |
| AQP9 | NM_020980 NM_001320635 NM_001320636 | AAGCTATTCTCAGTCGAGGAC | TCATCCGTCCAAAGAGACAC | 156 |
| FPR1 | NM_001193306 NM_002029 | TGCCAGTTATCATTCTGTGTG | AATGATGAACCGGATGATGC | 152 |
| FPR2 | NM_001462 NM_001005738 | TCTTTCACGGCCACATTACC | TTGATGTCCACCACGATGTG | 107 |
| NCF2 | NM_000433 NM_001127651 NM_001190794 NM_001190789 NM_001410895 | TCATGTTCAACGGGCAGAAG | CTTTGGAAGTAGGAGGAGCTG | 135 |

*bp: Base pair

Melt curve analysis (65-95 °C) verified product specificity (supplementary figures 2S-7S). All reactions were performed in technical duplicates. Relative gene expression data were analyzed using the $2^{-\Delta CT}$ [$2^{-(CT_{target\ gene}-CT_{reference\ gene})}$] method.²⁷

Statistical Analysis

The normality of the data distribution was evaluated using the Shapiro–Wilk test. For comparisons among more than two groups, the Kruskal–Wallis test was applied, followed by Dunn’s *post hoc* test with multiple-comparison correction. ROC curves were generated in GraphPad Prism (version 7.0; GraphPad Software, USA), and the optimal cutoff was determined using the Youden index. Data visualization was conducted with GraphPad Prism. Statistical significance was considered at P values less than 0.05, with non-significant results indicated as “ns”.

Results

Number of Participants

A total of 83 individuals were assessed for eligibility for the experimental validation phase of the study. Of these, 23 were excluded for not meeting the inclusion criteria, as 15 patients had a concurrent diagnosis of IBS, and eight patients had insufficient biopsy material. The final analysis included a total of 60 participants, comprising 20 healthy controls, 20 newly diagnosed UC patients, and 20 treatment-resistant UC patients.

Identification of DEGs in UC

The expression and distribution of all DEGs

with an adjusted P value of less than 0.01 and Log₂ fold change threshold =1 are illustrated in a volcano plot (figure 1). We identified 492 DEGs, which included 429 upregulated and 63 downregulated genes.




Figure 1: Volcano plot analysis of the GSE224758 dataset reveals 492 DEGs between UC patients and healthy controls. Red dots represent 429 significantly upregulated genes (adj. P<0.01, Log₂FC ≥1), while blue dots indicate 63 downregulated genes (adj. P<0.01, Log₂FC ≤-1). Black dots show non-significant genes. DEGs: Differentially expressed genes; UC: Ulcerative colitis.

PPI Network and Clustering

A co-expression PPI network for DEGs upregulated in UC compared to healthy controls was constructed using the STRING database. Clustering was performed using Cytoscape software with the Clusterviz plugin, revealing 14 modules (clusters). The largest module included 31 nodes and 83 edges, as shown in figure 2.




Figure 3: Pathway enrichment analysis of the largest gene module reveals significant enrichment of genes involved in the NETs formation pathway. Key genes highlighted include *FCGR3B*, *AQP9*, *FPR1*, *FPR2*, and *NCF2*. *FCGR3B*: Gamma Receptor IIIb; *AQP9*: Aquaporin-9; *FPR1*: Formyl Peptide Receptor 1; *FPR2*: Formyl Peptide Receptor 2; *NCF2*: Neutrophil cytosol factor 2.




Figure 4: Results of RT-qPCR analysis are shown, which was performed to quantify expression of five NET-related genes (*NCF2*, *FPR1*, *FPR2*, *AQP9*, and *FCGR3B*) in healthy controls (n=20), new UC cases (n=20), and treatment-resistant UC patients (n=20). Statistical analysis was performed using the Kruskal–Wallis test for overall group differences, followed by Dunn's *post hoc* test for pairwise comparisons. *NCF2*: No significant differences were observed among the three groups (Kruskal–Wallis $P=0.47$; all pairwise comparisons ns). *FPR1*: Expression was significantly increased in new UC cases compared with controls, while control vs. resistant and new case vs. resistant comparisons were non-significant. *FPR2*: Both new UC cases and resistant patients showed significantly higher expression relative to controls ($P=0.03$ and $P=0.02$, respectively), whereas new case vs. resistant comparisons were non-significant. *AQP9*: Expression was markedly elevated in both UC groups relative to controls (Kruskal–Wallis $P<0.0001$). Pairwise Mann–Whitney tests showed strong significance for control vs. new case ($P<0.0001$) and control vs. resistant ($P<0.0001$), while new case vs. resistant remained non-significant. Median values correspond to approximately 9–10-fold (new case) and 19–20-fold (resistant) increases compared with controls. *FCGR3B*: Expression trended higher in UC groups but did not reach statistical significance (Kruskal–Wallis $P=0.08$); all pairwise comparisons ns. Data are presented as individual values with horizontal bars indicating median±interquartile range. Significance levels are indicated in the figure as: ns: not significant; * $P<0.05$, **** $P<0.0001$. RT-qPCR: Reverse transcription-quantitative polymerase chain reaction; NET: Neutrophil Extracellular Traps; UC: Ulcerative colitis; *NCF2*: Neutrophil cytosol factor 2; *FPR1*: Formyl Peptide Receptor 1; *FPR2*: Formyl Peptide Receptor 2; *AQP9*: Aquaporin-9; *FCGR3B*: Gamma Receptor IIIb




Figure 5: ROC curve analysis of *AQP9* expression demonstrates its ability to discriminate UC patients from healthy controls. ROC analysis comparing newly diagnosed UC patients versus controls showed an AUC of 0.9382 (SE=0.04109; 95% CI: 0.8577-1.000; P<0.0001). ROC analysis comparing treatment-resistant UC patients versus controls yielded an AUC of 0.8789 (SE=0.06258; 95% CI: 0.7563-1.000; P<0.0001). Sensitivity and specificity values were calculated at the optimal cutoff determined using the Youden index in GraphPad Prism. These results indicate that *AQP9* exhibits strong discriminatory performance as a candidate tissue biomarker for UC. ROC: Receiver operating characteristic; AUC: area under the curve; UC: Ulcerative colitis; *AQP9*: Aquaporin-9; SE: Standard error

Consistently, ROC analysis comparing healthy controls and treatment-resistant UC patients showed strong discrimination, with an AUC of 0.8789 (SE=0.06258; 95% CI: 0.7563–1.000; P<0.0001). At the optimal cutoff, sensitivity was 80.98%, and specificity was 64.24%. Together, these findings indicate that *AQP9* expression exhibits high discriminatory accuracy for distinguishing UC patients from healthy individuals.

Discussion

This study set out to determine the key genes of NET formation in IBD patients compared to healthy individuals. Bioinformatic analysis led us to focus on five genes, including *FCGR3B*, *AQP9*, *FPR1*, *FPR2*, and *NCF2*. Our findings indicated that the *AQP9* gene was not only significantly upregulated in the new cases and the resistant patients compared with the control group, but it also showed a candidate tissue biomarker in UC patients.

AQP9 is an aquaglyceroporin expressed in immune cells, including neutrophils, where it contributes to water and solute transport processes that are important for inflammatory responses. Recent work in sepsis indicates that dysregulated aquaporin function, including *AQP9*, can modulate leukocyte recruitment and inflammatory injury, suggesting that altered *AQP9* activity may influence neutrophil-driven tissue damage in severe inflammation. Moreover, integrative transcriptomic analyses have shown that *AQP9* expression correlates with neutrophil infiltration and other immune signatures in diverse diseases, supporting its

role as an immune-related gene, although direct mechanistic evidence linking *AQP9* to NETosis in UC is still limited and warrants further investigation.^{28, 29}

In general, *AQP* genes are responsible for water channels, playing an essential role in maintaining water homeostasis in various tissues. They are strongly associated with IBD progression. Escudero and colleagues revealed that Aquaporin 8 (*AQP8*), as part of this family, was downregulated in collagenous colitis, while its expression was reversed upon corticosteroid therapy.³⁰ Consistently, Ricanek and colleagues showed that the expression of Aquaporin 3 (*AQP3*), 7, and 8 was reduced in IBD patients.³¹ Moreover, Bing Yu and others showed that *AQP9* expression was suppressed in Crohn's disease (CD), with a high diagnostic value.³² By contrast, Minhóa Yu and colleagues showed that in a refractory model of CD, *AQP9* expression was increased constantly. They demonstrated that anti-TNF therapy was more effective when complemented with inhibiting *AQP9*, whereby blocking *AQP9* decreased inflamed macrophage functions and cytokine expression, especially IL-23 and IL-1 β .³³ These findings, together with our data in UC, suggest that *AQP* expression patterns in IBD are highly context-dependent and tightly regulated to preserve intestinal homeostasis, and that disruption of this balance may contribute to disease onset and progression. Our results support the potential of *AQP9* for distinguishing UC patients from healthy individuals, while also underscoring the need for mechanistic studies to clarify how modulation of *AQP9* expression could be exploited therapeutically in UC.

FPR1 and *FPR2*, which play critical roles in neutrophil chemotaxis and chronic inflammation, were other genes involved in NET formation that were upregulated in patient groups. McAllister and others discovered that *FPR1* drives neutrophilic inflammation, leading to the development of IBD.³⁴ Furthermore, *FPR1* strongly contributes to the progression of CD by promoting macrophage M1 polarization.³⁵ Similarly, *FPR2* is highly expressed on neutrophils, where it drives their recruitment to inflamed colonic tissue. The interaction of *FPR2* with damage-associated molecular patterns (DAMPs) and with pathogen-associated molecular patterns (PAMPs) leads to neutrophil activation and NET formation. Implantation of NETs in the intestinal mucosa of IBD patients, with continued accumulation, results in functional epithelial barrier disruption, immune cell recruitment, and sustained inflammation.³⁶ This concept is also supported by our findings, considering the upregulation of *FPR2* in UC, which could contribute to exaggerated neutrophil-driven tissue damage. NET degradation defects have been implicated in IBD; therefore, *FPR2*-mediated NET may have a significant role in disease worsening. Collectively, it seems that co-expression of *FPR1* and *FPR2* causes IBD exacerbation through sustained hyperactivation of NET formation and macrophage M1 polarization. Therefore, targeting *FPR1* and *FPR2* could be a novel or complementary approach in the therapy for IBD.

NCF2 and *FCGR3B* were other candidate genes in the current study, and we did not find meaningful changes in their expression in the studied groups. There is little evidence to support *NCF2*'s contribution to IBD development. Muise and colleagues reported that a missense variant in *NCF2* might be associated with the onset of IBD.²⁵ There is also a weak association between *FCGR3B* and IBD, although the allele copy number of *FCGR3B* may have a significant association with UC susceptibility.³⁷ We suggest that NET formation is key to promoting both UC and CD, as supported by strong evidence, and *AQP9*, *FPR1*, and *FPR2* are critical in its formation. In the present study, we demonstrated that these three genes were upregulated in UC patients, suggesting these genes ensure the maintenance of activation of NET formation. We suggest inhibiting these genes as a complementary treatment alongside standard medical approaches for UC patients.

Several limitations must be acknowledged in this study. Firstly, smoking status, active infections, and medication use were not fully controlled, as these factors may affect mucosal gene expression profiles and potentially confound

biomarker discovery. Although patients with immune deficiencies were excluded, detailed records of smoking history and concurrent medications were not systematically collected due to constraints related to the sample size. Secondly, this cross-sectional design allows for the establishment of associations but does not prove causality between NET gene expression and the pathogenesis of UC. Thirdly, the relatively small sample size (n=60) limits the statistical power for subgroup analyses, particularly when comparing new and refractory UC patients. Finally, functional assays to confirm direct NET formation by the upregulated genes were not conducted. Future studies should involve larger cohorts, prospective designs, matched controls for confounding variables, and the validation of functional assays.

Conclusion

AQP9, *FPR1*, and *FPR2* are candidate tissue biomarkers for UC, with *AQP9* demonstrating excellent discriminatory performance. Further multi-center validation studies are necessary to confirm their clinical utility and establish their role in UC diagnosis.

Acknowledgment

The authors sincerely thank Kurdistan University of Medical Sciences. This study is extracted from the thesis of Hawra Zia Hossein al-Jamali (MSc), with grant number IR.MUK.REC.1403.285.

Authors' Contribution

H.A: Conceptualization, Funding acquisition, conducted experimental stages, data collection, and drafting; S.F: Conceptualization, study design, bioinformatics analyses, writing the final report, and drafting; M.M: Statistical analysis, manuscript writing and editing; A.A, A.J, and M.R.R: Sample collection, experimental stages, and reviewing the manuscript; B.N, as a pathologist, and F.S, as a gastroenterologist: Patient identification and editing the final draft of the manuscript. All authors have thoroughly reviewed and approved the final manuscript and agree to take full responsibility for all aspects of the work, ensuring that any questions concerning the accuracy or integrity of any part of the work are properly addressed.

Declaration of AI

The authors declare that no AI tools were used in the preparation of this manuscript.

Conflict of Interest: None declared.

References

- 1 Sosna B, Aebischer D, Myśliwiec A, Dynarowicz K, Bartusik-Aebischer D, Oleś P, et al. Selected Cytokines and Metalloproteinases in Inflammatory Bowel Disease. *Int J Mol Sci.* 2023;25. doi: 10.3390/ijms25010202. PubMed PMID: 38203373; PubMed Central PMCID: PMC10779120.
- 2 Tanwar P, Naagar M, Maity MK. A review on Ulcerative Colitis: Epidemiology, Risk Factors, Pathophysiology, Diagnosis, Clinical Management and Treatment Modalities. *International Journal of Enhanced Research in Medicines & Dental Care.* 2023;10:15-29.
- 3 Langan RC, Gotsch PB, Krafczyk MA, Skillinge DD. Ulcerative colitis: diagnosis and treatment. *Am Fam Physician.* 2007;76:1323-30. PubMed PMID: 18019875.
- 4 Burri E, Maillard MH, Schoepfer AM, Seibold F, Van Assche G, Rivière P, et al. Treatment Algorithm for Mild and Moderate-to-Severe Ulcerative Colitis: An Update. *Digestion.* 2020;101:2-15. doi: 10.1159/000504092. PubMed PMID: 31945767.
- 5 van Lierop PP, Samsom JN, Escher JC, Nieuwenhuis EE. Role of the innate immune system in the pathogenesis of inflammatory bowel disease. *J Pediatr Gastroenterol Nutr.* 2009;48:142-51. doi: 10.1097/MPG.0b013e3181821964. PubMed PMID: 19179875.
- 6 Shen ZH, Zhu CX, Quan YS, Yang ZY, Wu S, Luo WW, et al. Relationship between intestinal microbiota and ulcerative colitis: Mechanisms and clinical application of probiotics and fecal microbiota transplantation. *World J Gastroenterol.* 2018;24:5-14. doi: 10.3748/wjg.v24.i1.5. PubMed PMID: 29358877; PubMed Central PMCID: PMC5757125.
- 7 Zhou G, Yu L, Fang L, Yang W, Yu T, Miao Y, et al. CD177(+) neutrophils as functionally activated neutrophils negatively regulate IBD. *Gut.* 2018;67:1052-63. doi: 10.1136/gutjnl-2016-313535. PubMed PMID: 28468761.
- 8 Fukunaga S, Kuwaki K, Mitsuyama K, Takedatsu H, Yoshioka S, Yamasaki H, et al. Detection of calprotectin in inflammatory bowel disease: Fecal and serum levels and immunohistochemical localization. *Int J Mol Med.* 2018;41:107-18. doi: 10.3892/ijmm.2017.3244. PubMed PMID: 29115397; PubMed Central PMCID: PMC5746327.
- 9 Drury B, Hardisty G, Gray RD, Ho GT. Neutrophil Extracellular Traps in Inflammatory Bowel Disease: Pathogenic Mechanisms and Clinical Translation. *Cell Mol Gastroenterol Hepatol.* 2021;12:321-33. doi: 10.1016/j.jcmgh.2021.03.002. PubMed PMID: 33689803; PubMed Central PMCID: PMC8166923.
- 10 Long D, Mao C, Xu Y, Zhu Y. The emerging role of neutrophil extracellular traps in ulcerative colitis. *Front Immunol.* 2024;15:1425251. doi: 10.3389/fimmu.2024.1425251. PubMed PMID: 39170617; PubMed Central PMCID: PMC11335521.
- 11 Josefs T, Barrett TJ, Brown EJ, Quezada A, Wu X, Voisin M, et al. Neutrophil extracellular traps promote macrophage inflammation and impair atherosclerosis resolution in diabetic mice. *JCI Insight.* 2020;5. doi: 10.1172/jci.insight.134796. PubMed PMID: 32191637; PubMed Central PMCID: PMC7205252.
- 12 Kim TS, Silva LM, Theofilou VI, Greenwell-Wild T, Li L, Williams DW, et al. Neutrophil extracellular traps and extracellular histones potentiate IL-17 inflammation in periodontitis. *J Exp Med.* 2023;220. doi: 10.1084/jem.20221751. PubMed PMID: 37261457; PubMed Central PMCID: PMC10236943.
- 13 Zhang Y, Guo L, Dai Q, Shang B, Xiao T, Di X, et al. A signature for pan-cancer prognosis based on neutrophil extracellular traps. *J Immunother Cancer.* 2022;10. doi: 10.1136/jitc-2021-004210. PubMed PMID: 35688556; PubMed Central PMCID: PMC9189842.
- 14 Li J, Liu Y, Sun Z, Zeng S, Zheng C. Identification of neutrophil extracellular trap-related biomarkers in ulcerative colitis based on bioinformatics and machine learning. *Front Genet.* 2025;16:1589999. doi: 10.3389/fgene.2025.1589999. PubMed PMID: 40620701; PubMed Central PMCID: PMC12226468.
- 15 Xv Y, Feng Y, Lin J. CXCR1 and CXCR2 are potential neutrophil extracellular trap-related treatment targets in ulcerative colitis: insights from Mendelian randomization, colocalization and transcriptomic analysis. *Front Immunol.* 2024;15:1425363. doi: 10.3389/fimmu.2024.1425363. PubMed PMID: 39328405; PubMed Central PMCID: PMC11424450.
- 16 Ou J, Li L, Zhi F, Huang B, Zhao X. Developing models for the diagnosing of ulcerative colitis and prognosis of anti-TNF- α non-response based on neutrophil extracellular trap-associated genes. *Front Immunol.* 2025;16:1530508. doi: 10.3389/fimmu.2025.1530508. PubMed PMID: 40963602; PubMed Central PMCID: PMC12436428.
- 17 Alemán OR, Mora N, Cortes-Vieyra R,

- Uribe-Querol E, Rosales C. Differential Use of Human Neutrophil Fcγ Receptors for Inducing Neutrophil Extracellular Trap Formation. *J Immunol Res.* 2016;2016:2908034. doi: 10.1155/2016/2908034. PubMed PMID: 27034964; PubMed Central PMCID: PMC4806689.
- 18 Liang P, Zhu M, Sun X, Wang L, Li B, Ming S, et al. LncRNA-mRNA co-expression analysis reveals aquaporin-9-promoted neutrophil extracellular trap formation and inflammatory activation in sepsis. *Int Immunopharmacol.* 2024;140:112916. doi: 10.1016/j.intimp.2024.112916. PubMed PMID: 39133961.
- 19 da Silva IV, Garra S, Calamita G, Soveral G. The Multifaceted Role of Aquaporin-9 in Health and Its Potential as a Clinical Biomarker. *Biomolecules.* 2022;12. doi: 10.3390/biom12070897. PubMed PMID: 35883453; PubMed Central PMCID: PMC9313442.
- 20 Nafiz TN, Sankar P, Mishra LK, Rousseau RP, Saqib M, Subbian S, et al. Differential requirement of formyl peptide receptor 1 in macrophages and neutrophils in the host defense against *Mycobacterium tuberculosis* Infection. *Sci Rep.* 2024;14:23595. doi: 10.1038/s41598-024-71180-1. PubMed PMID: 39384825; PubMed Central PMCID: PMC11464745.
- 21 Lou X, Chen H, Chen S, Ji H, He T, Chen H, et al. LL37/FPR2 regulates neutrophil mPTP promoting the development of neutrophil extracellular traps in diabetic retinopathy. *Faseb j.* 2024;38:e23697. doi: 10.1096/fj.202400656R. PubMed PMID: 38842874.
- 22 Yi X, Tran E, Odiba JO, Qin CX, Ritchie RH, Baell JB. The formyl peptide receptors FPR1 and FPR2 as targets for inflammatory disorders: recent advances in the development of small-molecule agonists. *Eur J Med Chem.* 2024;265:115989. doi: 10.1016/j.ejmech.2023.115989. PubMed PMID: 38199163.
- 23 Song Z, Yoo DG, Idol RA, Barbu EA, Jacob CO, Dinauer MC. Lupus-associated NCF2 variant p.R395W in the NADPH oxidase 2 complex results in a reduced production of reactive oxygen species by myeloid cells. *Front Lupus.* 2023;1:1186641. doi: 10.3389/flu.2023.1186641.
- 24 Garley M, Dziemiańczyk-Pakieła D, Ratajczak-Wrona W, Prczynicz A, Nowak K, Łazarczyk B, et al. NETs biomarkers in saliva and serum OSCC patients: One hypothesis, two conclusions. *Adv Med Sci.* 2022;67:45-54. doi: 10.1016/j.advms.2021.12.004. PubMed PMID: 34971930.
- 25 Muise AM, Xu W, Guo CH, Walters TD, Wolters VM, Fattouh R, et al. NADPH oxidase complex and IBD candidate gene studies: identification of a rare variant in NCF2 that results in reduced binding to RAC2. *Gut.* 2012;61:1028-35. doi: 10.1136/gutjnl-2011-300078. PubMed PMID: 21900546; PubMed Central PMCID: PMC3806486.
- 26 Fakhari S, Moradzad M, Ahmadi A, Hekmatnia M, Jalili A, Nikkhoo B, et al. Down-regulation of MT2-MMP and MT5-MMP in ulcerative colitis serves a diagnostic predictor and potential therapeutic targets. *Mol Biol Rep.* 2025;52:359. doi: 10.1007/s11033-025-10449-8. PubMed PMID: 40178713.
- 27 Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc.* 2008;3:1101-8. doi: 10.1038/nprot.2008.73. PubMed PMID: 18546601.
- 28 Rump K, Adamzik M. Aquaporins in sepsis-an update. *Front Immunol.* 2024;15:1495206. doi: 10.3389/fimmu.2024.1495206. PubMed PMID: 39544938; PubMed Central PMCID: PMC11560437.
- 29 Liu X, Xu Q, Li Z, Xiong B. Integrated analysis identifies AQP9 correlates with immune infiltration and acts as a prognosticator in multiple cancers. *Sci Rep.* 2020;10:20795. doi: 10.1038/s41598-020-77657-z. PubMed PMID: 33247170; PubMed Central PMCID: PMC7699650.
- 30 Escudero-Hernández C, Münch A, Østvik AE, Granlund AVB, Koch S. The Water Channel Aquaporin 8 is a Critical Regulator of Intestinal Fluid Homeostasis in Collagenous Colitis. *J Crohns Colitis.* 2020;14:962-73. doi: 10.1093/ecco-jcc/jjaa020. PubMed PMID: 32016376; PubMed Central PMCID: PMC7393183.
- 31 Ricanek P, Lunde LK, Frye SA, Støen M, Nygård S, Morth JP, et al. Reduced expression of aquaporins in human intestinal mucosa in early stage inflammatory bowel disease. *Clin Exp Gastroenterol.* 2015;8:49-67. doi: 10.2147/ceg.s70119. PubMed PMID: 25624769; PubMed Central PMCID: PMC4296881.
- 32 Yu B, Yin YX, Tang YP, Wei KL, Pan ZG, Li KZ, et al. Diagnostic and Predictive Value of Immune-Related Genes in Crohn's Disease. *Front Immunol.* 2021;12:643036. doi: 10.3389/fimmu.2021.643036. PubMed PMID: 33936061; PubMed Central PMCID: PMC8085323.
- 33 Yu M, Shi Y, Gao Y, Luo Y, Jin Y, Liang X, et al. Targeting AQP9 enhanced the anti-TNF therapy response in Crohn's disease by inhibiting LPA-hippo pathway. *Pharmacol*

- Res. 2024;203:107172. doi: 10.1016/j.phrs.2024.107172. PubMed PMID: 38583685.
- 34 McAllister MJ, Hall R, Whelan RJ, Fischer LJ, Chuah CS, Cartlidge PD, et al. Formylated Peptide Receptor-1-Mediated Gut Inflammation as a Therapeutic Target in Inflammatory Bowel Disease. *Crohns Colitis* 360. 2024;6:otae003. doi: 10.1093/crocol/otae003. PubMed PMID: 38352118; PubMed Central PMCID: PMC10862654.
- 35 Ye C, Zhu S, Yuan J, Yuan X. FPR1, as a Potential Biomarker of Diagnosis and Infliximab Therapy Responses for Crohn's Disease, is Related to Disease Activity, Inflammation and Macrophage Polarization. *J Inflamm Res*. 2024;17:3949-66. doi: 10.2147/jir.s459819. PubMed PMID: 38911989; PubMed Central PMCID: PMC11193993.
- 36 Yang WS, Wang JL, Wu W, Wang GF, Yan J, Liu Q, et al. Formyl peptide receptor 2 as a potential therapeutic target for inflammatory bowel disease. *Acta Pharmacol Sin*. 2023;44:19-31. doi: 10.1038/s41401-022-00944-0. PubMed PMID: 35840658; PubMed Central PMCID: PMC9812994.
- 37 Asano K, Matsumoto T, Umeno J, Hirano A, Esaki M, Hosono N, et al. Impact of allele copy number of polymorphisms in FCGR3A and FCGR3B genes on susceptibility to ulcerative colitis. *Inflamm Bowel Dis*. 2013;19:2061-8. doi: 10.1097/MIB.0b013e318298118e. PubMed PMID: 23917248.