

The Gut–Brain Axis in Sleep Deprivation: Mechanisms, Dysbiosis, and Potential Therapeutic Avenues

Esmail Karami¹, PhD;^{ORCID} Mohammad Taghi Mohammadi², PhD; Hossein Shirvani¹, PhD; Behzad Bazgir¹, PhD^{ORCID}

¹Exercise Physiology Research Center, Life Style Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran;

²Department of Physiology and Medical Physics, Faculty of Medicine, Baqiyatallah University of Medical Sciences, Tehran, Iran

Correspondence:

Behzad Bazgir, PhD;
Baqiyatallah University of Medical Sciences, Vanak Sq., Mulla Sadra St., Sheikh Bahai St., Postal code: 14155-64371, Tehran, Iran

Tel: +98 21 88754475

Email: bazgirbehzad@yahoo.com

Received: 13 September 2025

Revised: 25 October 2025

Accepted: 16 November 2025

Abstract

Sleep is fundamental to metabolic regulation, cardiovascular health, immune balance, and cognitive performance. Emerging evidence indicates that sleep deprivation alters gut microbial communities and immune signaling, thereby influencing the gut-brain axis. This narrative review synthesized current findings on how sleep loss engages gut–brain pathways and evaluated therapeutic avenues linked to specific mechanisms, drawing on peer-reviewed human and animal studies and prioritizing sleep- and brain-relevant outcomes. Evidence converged on three mechanistic pillars: (1) circadian and microbial misalignment, which disrupts diurnal microbiome dynamics; (2) inflammatory and barrier dysfunction loops that increase permeability and immune activation; and (3) altered neurotransmitter and metabolite signaling, including shifts in tryptophan pathways and reduced short-chain fatty acid production, with downstream effects on neuroinflammation and plasticity. Human data supported associations between sleep loss, poor sleep quality, and impaired cognition. However, causal inference was limited by small sample sizes, study heterogeneity, and mixed endpoints. Interventions such as probiotics, prebiotics, postbiotics, synbiotics, diet-based short-chain fatty acid strategies, and chronobiotics are biologically plausible, yet they are supported mainly by small trials or preclinical models. Fecal microbiota transplantation shows exploratory, yet faces significant safety, standardization, and regulatory challenges. Collectively, the evidence indicated that sleep deprivation disrupts gut microbiota composition and gut-brain signaling, with consequences for neuroinflammation, neurotransmitter balance, and cognition. This review highlighted the consequent need for standardized endpoints, strain-resolved microbiome analyses, and well-powered randomized controlled trials.

Please cite this article as: Karami E, Mohammadi MT, Shirvani H, Bazgir B. The Gut–Brain Axis in Sleep Deprivation: Mechanisms, Dysbiosis, and Potential Therapeutic Avenues. *Iran J Med Sci*. 2026;51(8):533-542. doi: 10.30476/ijms.2025.108475.4351.

Keywords • Sleep deprivation • Gut–brain axis (GBA) • Microbiota • Neuroinflammation • Probiotics

What's Known

- Insufficient sleep is associated with metabolic, cardiovascular, and mental health risks.
- Sleep loss perturbs gut microbial communities and immune signaling, which can influence brain function.

What's New

- We propose a tiered model illustrating how sleep loss disrupts the gut–brain axis through (i) circadian and microbial misalignment, (ii) inflammatory and barrier dysfunction loops, and (iii) altered neurotransmitter and metabolite signaling.
- We mapped interventions to these specific mechanisms, including nutritional strategies targeting short-chain fatty acids, postbiotics, microbiota-informed chronobiotics, probiotics, and fecal microbiota transplantation.

Introduction

Sleep deprivation has become a widespread problem in contemporary life, exerting a substantial negative influence on productivity and general well-being across diverse populations.¹ Chronic sleep deprivation is linked to several major health issues, such as metabolic disorders, cardiovascular diseases, and mental health impairments, in addition to the obvious ones,

such as impaired cognitive performance and excessive daytime sleepiness.² The established connection between sleep deprivation and brain or mental health issues underscores the urgency of addressing this public health challenge.³

According to recent research, sleep deprivation has a significant impact on the gut microbiota, which is the intricate ecosystem of microorganisms residing in the gastrointestinal tract.⁴ Sleep disturbances can cause dysbiosis, characterized by an imbalance between beneficial and harmful bacteria and a reduction in microbial diversity.⁵ Numerous factors, such as changes in circadian rhythms, dietary practices, and stress hormone levels, are thought to be responsible for these changes in the composition of the gut microbiota. Since many physiological processes depend on the gut microbiota, its disruption can have cascading effects on overall health.⁶⁻⁸

Notably, the gut-brain axis (GBA) serves as the bidirectional communication network linking the gut and the brain, influencing emotion, cognition, and behavior.⁹ Dysbiosis induced by sleep deprivation may disrupt GBA signaling, potentially exacerbating neurological dysfunction.¹⁰ This complex relationship emphasizes the critical need to understand the specific mechanisms by which sleep deprivation impairs gut-brain communication.¹¹

This narrative review aimed to investigate the effects of sleep deprivation on gut-brain crosstalk and to evaluate the potential of emerging interventions designed to mitigate these detrimental effects. By synthesizing the current literature, this review aimed to provide a mechanistic foundation for future research and therapeutic strategies aimed at preserving gut and brain health in the context of sleep loss.

Literature Selection

This narrative review integrated multidisciplinary evidence on how sleep deprivation alters the gut-brain axis, prioritizing mechanistic and translational synthesis over quantitative meta-analysis. A structured search of MEDLINE (via PubMed) and Scopus was conducted for literature published up to June 2025. The search strategy was designed to identify studies on sleep loss, microbiome dynamics, inflammation, neurotransmitters, short-chain fatty acids (SCFAs), and fecal microbiota transplantation (FMT). Relevant studies were also identified through hand-searching of reference lists.

Eligible studies included peer-reviewed, English-language reports of human, animal, and *in vitro* studies that specifically addressed

microbiota composition, microbial metabolites, intestinal barrier and immune integrity, neurological outcomes, or microbiota-directed interventions. Non-English reports, conference abstracts, and opinion pieces were excluded. Data pertaining to study design, populations, outcomes, and mechanisms were extracted and narratively synthesized. The synthesis was organized around three focal domains: (i) circadian microbial misalignment, (ii) inflammatory and barrier dysfunction, and (iii) neurotransmitter and metabolite signaling. This structure provided an integrative framework for linking sleep deprivation-induced dysbiosis to neuroinflammation and cognitive impairment.

Effects of Sleep Deprivation on the Gut Microbiota

As discussed, sleep deprivation significantly disrupts the balance of the gut microbiota, leading to substantial alterations in its diversity and composition. It is associated not only with a general decline in microbial richness but also with specific changes in the relative abundance of specific bacterial species.^{12, 13}

A primary cause of this disruption is the disturbance of circadian rhythms, which are crucial for regulating the diurnal oscillations of gut bacteria.¹⁴ Circadian rhythms coordinate the timing of microbial growth and activity, which is why beneficial microorganisms thrive during wakefulness and digestion periods.¹⁵ This timing is disrupted when sleep is interrupted, which can result in changes in intestinal motility and nutrient absorption, thereby creating an environment conducive to dysbiosis.^{16, 17}

Furthermore, sleep deprivation activates the hypothalamic-pituitary-adrenal axis, elevating levels of stress hormones such as cortisol.¹⁸ These hormones directly impact the gut environment, promoting the proliferation of stress-resistant species while suppressing beneficial ones. This modification establishes a deleterious feedback loop: the altered microbiome exacerbates local inflammation, which in turn further compromises gut barrier integrity.^{6, 19, 20}

Compounding the issue, sleep deprivation induces dysregulation of the immune system, impairing gut mucosal immunity. In addition to exacerbating dysbiosis, the subsequent immune dysfunction feeds systemic inflammation, which can subsequently affect the central nervous system (CNS) and other organ systems.²¹

Dietary patterns are also altered by inadequate sleep, frequently resulting in increased consumption of high-fat and high-calorie foods.^{22, 23} These dietary modifications further influence the composition and metabolic activity of the gut

microbiota by providing substrates that promote the growth of specific microbial populations. For example, a diet high in processed foods can favor the proliferation of bacteria that thrive on simple sugars and fats, while depleting beneficial microbes dependent on dietary fiber. This imbalance can contribute to metabolic disorders such as insulin resistance and obesity, which promote inflammation detrimental to gastrointestinal health.²⁴

Additionally, inadequate sleep can induce epigenetic modifications in both the gut microbiota and host cells. These alterations in gene expression can influence microbial metabolic activity and the overall gut ecosystem. Such epigenetic modifications have the potential to further complicate the relationship between sleep, diet, and gut health by modulating how gut bacteria interact with the host and metabolize nutrients.^{25, 26}

Of note, the systemic effects of sleep deprivation extend beyond the intestine, substantially impacting the production of gut microbial metabolites, particularly SCFAs.²⁷ SCFAs, including butyrate, acetate, and propionate, are primarily produced through the bacterial fermentation of indigestible dietary fiber.²⁸ These metabolites are essential for maintaining gastrointestinal health and influence diverse physiological processes, including brain function.²⁹

A critical role of SCFAs is to regulate the gut barrier integrity. They serve as a primary energy source for colonocytes and enhance the expression of tight junction proteins. However, sleep deprivation disrupts SCFA synthesis, leading to a decrease in the concentration of these beneficial metabolites.³⁰⁻³² This reduction can compromise intestinal barrier integrity and impair immunological responses.^{33, 34}

As SCFAs can cross the blood-brain barrier (BBB), their altered production has significant implications for brain health. Reduced levels of butyrate, for instance, may exacerbate neuroinflammation by failing to inhibit the production of pro-inflammatory cytokines, thereby increasing susceptibility to cognitive impairment, mood disorders, and neurodegenerative diseases.^{35, 36} Additionally, SCFAs influence the expression of brain-derived neurotrophic factor (BDNF), a key regulator of synaptic plasticity and neurogenesis, directly linking gut metabolic output to brain function and resilience.³⁷

Therapeutic Implications and Research Priorities

This narrative review provided an integrative synthesis of the current evidence linking sleep

deprivation to GBA dysfunction. It emphasized the mechanistic interplay between circadian misalignment, microbial dysbiosis, immune–barrier disruption, and altered neurochemical signaling. Rather than performing a quantitative meta-analysis, this review critically consolidated mechanistic insights from human and animal studies to delineate how chronic sleep loss could reprogram gut microbiota composition, disrupt metabolite and neurotransmitter pathways, and ultimately impair cognitive and emotional regulation.³⁸⁻⁴⁴

The analysis identified three interconnected mechanistic domains through which sleep deprivation exerts its deleterious effects: (i) the disruption of circadian–microbial synchronization, (ii) the induction of inflammatory and epithelial barrier dysfunction loops, and (iii) the alteration of neurotransmitter and metabolite signaling, particularly involving tryptophan derivatives and SCFAs.^{27, 28, 32, 35-37} Collectively, these findings suggest that gut microbial imbalance serves as both a mediator and amplifier of sleep-related neurophysiological disturbances.^{21, 39, 40, 43, 44}

By integrating these mechanisms, the review advanced a conceptual understanding of gut–brain cross-talk in sleep biology and highlighted several biologically plausible therapeutic strategies, such as probiotics, prebiotics, and FMT, that may restore microbial homeostasis and mitigate neuroinflammatory consequences of sleep deprivation.⁴⁵⁻⁵⁰

The Promising Role of Newly Emerged Interventions

Building on these mechanistic insights, recent studies have explored potential interventions targeting the gut–brain axis to mitigate the effects of sleep deprivation. Clinical and preclinical data suggest that modulating the gut microbiota can have beneficial effects on mood, cognition, and sleep regulation.⁵¹⁻⁵³ Probiotics and prebiotics have demonstrated potential in restoring microbial balance and enhancing the production of beneficial metabolites, such as SCFAs, which are known to strengthen intestinal barrier integrity and reduce systemic inflammation.^{46, 48}

The FMT has emerged as a particularly promising, though still experimental, intervention. Studies indicated that FMT can improve sleep quality and neurocognitive outcomes by correcting dysbiosis and enhancing microbial diversity.^{45, 50} In patients with chronic fatigue or sleep-related cognitive impairment, FMT interventions was shown to partially restore a healthy microbial composition and alleviate neuroinflammatory symptoms.^{47, 49}

Despite its promise, FMT faces significant safety, standardization, and regulatory challenges that currently limit its broad clinical application.⁴⁹ Collectively, this converging evidence underscores the translational potential of microbiota-targeted therapies for addressing sleep deprivation-related dysfunctions.⁴⁵⁻⁵³

These findings collectively underscored the complex interplay between sleep deprivation, the gut microbiota, and neurological outcomes, and pointed to probiotics, prebiotics, and other microbiota-focused strategies as promising therapeutic avenues. A summary of the main findings is presented in table 1.

Integrative Conceptual Framework: Linking Gut Dysbiosis to Neuroinflammation and Cognitive Impairment

To move beyond descriptive reporting, this current synthesis proposes an integrative conceptual framework that illustrates how sleep deprivation may causally perturb gut-brain communication. Chronic sleep loss disrupts the circadian regulation of gut microbial oscillations, leading to dysbiosis characterized by a loss of beneficial taxa (e.g., *Lactobacillus*, *Bifidobacterium*) and reduced microbial diversity.^{12, 13, 16} This imbalance increases intestinal permeability ("leaky gut"), allowing translocation of

bacterial components such as lipopolysaccharides (LPS) into systemic circulation.^{43, 44}

The resulting systemic and neuroinflammation activates immune-neuroendocrine cascades, particularly via the HPA axis, elevating levels of cortisol and pro-inflammatory cytokines (e.g., IL-6, TNF- α).^{39, 40, 42} Concurrently, dysbiosis alters the metabolism of tryptophan and SCFAs, reducing the production of serotonin and butyrate—metabolites that are crucial for neuronal plasticity and mood regulation.^{27, 28, 30, 32, 37, 54, 55}

Several studies demonstrated that this disruption extends to neurotransmitter synthesis and signaling, particularly serotonin, dopamine, and γ -aminobutyric acid (GABA) pathways. Sleep deprivation impairs tryptophan metabolism, reducing serotonin synthesis in both gut and brain, which contributes to mood instability and anxiety.⁵⁴⁻⁵⁸ Dopaminergic signaling is also affected; gut-derived dopamine precursors are reduced during dysbiosis, thereby altering reward and cognitive processing circuits.⁵⁹⁻⁶⁴ Similarly, populations of GABA-producing microbes, such as *Lactobacillus* and *Bifidobacterium*, are diminished by sleep deprivation, which impairs inhibitory neurotransmission and sleep quality.⁶⁵⁻⁶⁹

Collectively, these multi-level alterations; circadian dysregulation, barrier breakdown, immune activation, and neurochemical

Table 1: Summary of main findings

Main context	Main findings	References
Disruption of gut microbiota by sleep deprivation	Sleep deprivation leads to reduced diversity and altered bacteria; circadian rhythms are crucial for gut health.	12-14
Impact on gut health and dysbiosis	Disrupted sleep negatively affects intestinal motility and nutrient absorption, promoting dysbiosis and harming beneficial bacteria.	16-18
Dietary changes due to sleep deprivation	Sleep loss alters dietary habits, increasing the consumption of high-fat and calorie-dense foods, which fosters harmful gut microbes and metabolic disorders.	22-24
Epigenetic changes in gut microbiota	Sleep deprivation causes epigenetic modifications that impact microbial gene expression, complicating the interplay between sleep, diet, and gut health.	25, 26
Effects of gut microbial metabolites	Reduced production of SCFAs leads to neuroinflammation and cognitive decline.	28
Neurotransmitter production and sleep	Disruption affects tryptophan metabolism and neurotransmitters (serotonin, GABA), contributing to mood disorders.	54, 55
Inflammation and gut-brain crosstalk	Inflammatory molecules from gut bacteria contribute to systemic inflammation, increasing risks for neurological conditions such as Alzheimer's and depression.	39, 40, 43, 44
Experimental evidence of gut microbial changes	Animal studies show increased Firmicutes and decreased Bacteroidetes linked to colonic inflammation and cognitive impairments.	38, 42, 43
Clinical evidence on sleep deprivation and neurological outcomes	Sleep deprivation is linked to cognitive decline and mood disorders; specific microbiota are associated with cognition.	51-53
Gut microbiota and mental health	Insomnia, lower diversity, and mood disorder links; FMT improves sleep quality and mood.	45, 50
Interventional studies linking sleep and gut health	FMT may mitigate sleep deprivation effects, potential for chronic fatigue syndrome treatment.	47, 49
Probiotics and cognitive decline in older adults	Poor sleep quality in older adults is linked to cognitive decline; probiotics are a potential intervention.	46, 48

SCFAs: Short-chain fatty acids; GABA: γ -Aminobutyric acid; FMT: Fecal microbiota transplant

imbalance; disrupt the balance between excitatory and inhibitory neurotransmission, sustain a state of neuroinflammation, and ultimately exacerbate cognitive and emotional deficits.^{38, 41, 70, 71}

Therapeutic Implications

The proposed framework underscores a bidirectional, self-perpetuating cycle: sleep deprivation induces dysbiosis and systemic inflammation, which in turn worsen sleep architecture and cognitive function.^{39, 40, 43, 44} Breaking this cycle through microbiota-targeted interventions, such as probiotics, prebiotics, SCFA-enhancing diets, and FMT, offers a promising mechanistic strategy.⁴⁵⁻⁵⁰

Future research should prioritize longitudinal, mechanism-anchored randomized controlled trials (RCTs) that integrate multi-omic profiling (e.g., metagenomics, metabolomics) with neuroimaging to validate these proposed causal pathways and intervention efficacy.^{47, 49, 52}

Policy Implications and Future Directions

The findings of this review underscore the urgent need for public health policies that prioritize sleep health as a critical determinant of physical and mental well-being. Public awareness programs should highlight the established links between chronic sleep deprivation, gut dysbiosis, and cognitive or emotional disorders.^{21, 41} Preventive health strategies that incorporate sleep hygiene education, dietary regulation, and microbiota-supportive nutrition could play a pivotal role in reducing the societal burden of sleep-related illnesses.^{7, 24, 27}

Future research must prioritize longitudinal and multi-omics clinical trials to evaluate how targeted microbiome interventions, such as specific probiotic strains, prebiotic formulations, or FMT, can effectively restore gut-brain homeostasis.^{47, 49, 50, 52} Collaborative efforts between clinicians, neuroscientists, and microbiome researchers are essential to translate these mechanistic findings into actionable therapeutic guidelines and informed public health frameworks.^{46, 48}

Such integrative, translational approaches will not only deepen our understanding of the gut-brain-sleep triad but also guide the development of evidence-based strategies to improve sleep and neurological health at a population level.

Conclusion

In conclusion, this narrative review synthesized

evidence establishing that sleep deprivation induces gut dysbiosis, which subsequently disrupts the gut-brain axis, primarily through increased systemic inflammation and the dysregulation of key neurotransmitters. This mechanism ultimately contributes to cognitive deficits and mood disorders. The collective findings underscored the critical public health importance of addressing sleep deprivation. Furthermore, they indicated that microbiota-targeted interventions, such as specific probiotic formulations and FMT, represent a promising therapeutic avenue for restoring gut-brain homeostasis and improving neurological outcomes in affected individuals.

Acknowledgment

I would like to sincerely thank all the professors who contributed to this project for their valuable guidance, insightful comments, and continuous support throughout the development of this review.

Authors' Contribution

E.K and B.B were involved in all stages of the review, including conceptualization, literature search, analysis, and manuscript drafting. H.S and M.T.M contributed to the literature review, manuscript drafting, and critical revision. All authors have read and approved the final manuscript and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Declaration of AI

During the preparation of this manuscript, PaperGuide AI (web-based platform) was used to assist with literature searches, identification of relevant references, and organization of the bibliography. ChatGPT (OpenAI; GPT-4o) was used to improve the language, grammar, clarity, and overall academic writing style of the manuscript. AI tools were used solely to support these tasks and were not involved in generating scientific data, analyzing results, interpreting findings, or drawing the study's conclusions. All AI-assisted content was carefully reviewed, verified, and edited by the authors, who accept full responsibility for the accuracy, integrity, and final content of the manuscript.

Conflict of Interest: None declared.

References

- Guo C, Piao S, Wang C, Yu L, Wang K, Qu Q, et al. The prevalence and associated factors of sleep deprivation among healthy college students in China: a cross-sectional survey. *PeerJ*. 2023;11:e16009. doi: 10.7717/peerj.16009. PubMed PMID: 37744238; PubMed Central PMCID: PMC10512935.
- Liew SC, Aung T. Sleep deprivation and its association with diseases- a review. *Sleep Med*. 2021;77:192-204. doi: 10.1016/j.sleep.2020.07.048. PubMed PMID: 32951993.
- Mukherjee U, Sehar U, Brownell M, Reddy PH. Mechanisms, consequences and role of interventions for sleep deprivation: Focus on mild cognitive impairment and Alzheimer's disease in elderly. *Ageing Res Rev*. 2024;100:102457. doi: 10.1016/j.arr.2024.102457. PubMed PMID: 39154978; PubMed Central PMCID: PMC11415889.
- Smith RP, Easson C, Lyle SM, Kapoor R, Donnelly CP, Davidson EJ, et al. Gut microbiome diversity is associated with sleep physiology in humans. *PLoS One*. 2019;14:e0222394. doi: 10.1371/journal.pone.0222394. PubMed PMID: 31589627; PubMed Central PMCID: PMC6779243.
- Sun J, Fang D, Wang Z, Liu Y. Sleep Deprivation and Gut Microbiota Dysbiosis: Current Understandings and Implications. *Int J Mol Sci*. 2023;24. doi: 10.3390/ijms24119603. PubMed PMID: 37298553; PubMed Central PMCID: PMC10253795.
- Beurel E. Stress in the microbiome-immune crosstalk. *Gut Microbes*. 2024;16:2327409. doi: 10.1080/19490976.2024.2327409. PubMed PMID: 38488630; PubMed Central PMCID: PMC10950285.
- Singh RK, Chang HW, Yan D, Lee KM, Ucmak D, Wong K, et al. Influence of diet on the gut microbiome and implications for human health. *J Transl Med*. 2017;15:73. doi: 10.1186/s12967-017-1175-y. PubMed PMID: 28388917; PubMed Central PMCID: PMC5385025.
- Tian Y, Yang W, Chen G, Men C, Gu Y, Song X, et al. An important link between the gut microbiota and the circadian rhythm: imply for treatments of circadian rhythm sleep disorder. *Food Sci Biotechnol*. 2022;31:155-64. doi: 10.1007/s10068-021-01015-6. PubMed PMID: 35186346; PubMed Central PMCID: PMC8817960.
- Carabotti M, Scirocco A, Maselli MA, Severi C. The gut-brain axis: interactions between enteric microbiota, central and enteric nervous systems. *Ann Gastroenterol*. 2015;28:203-9. PubMed PMID: 25830558; PubMed Central PMCID: PMC4367209.
- Ullah H, Arbab S, Tian Y, Liu CQ, Chen Y, Qijie L, et al. The gut microbiota-brain axis in neurological disorder. *Front Neurosci*. 2023;17:1225875. doi: 10.3389/fnins.2023.1225875. PubMed PMID: 37600019; PubMed Central PMCID: PMC10436500.
- Wang Z, Wang Z, Lu T, Chen W, Yan W, Yuan K, et al. The microbiota-gut-brain axis in sleep disorders. *Sleep Med Rev*. 2022;65:101691. doi: 10.1016/j.smrv.2022.101691. PubMed PMID: 36099873.
- Das M, Muralitharan G, Saini S, Patra S. Sleep, gut microbiota, and mind-body medicine. *Brain Behavior and Immunity Integrative*. 2025;11:100128. doi: 10.1016/j.bbii.2025.100128.
- Supasitdikul T, Mazariegos JRR, Nhat NN, Tung YT, Yang DF, Lee LJ, et al. Sleep Deprivation Alters Gut Microbiome Diversity and Taxonomy: A Systematic Review and Meta-Analysis of Human and Rodent Studies. *J Sleep Res*. 2026;35:e70125. doi: 10.1111/jsr.70125. PubMed PMID: 40562421.
- Lotti S, Dinu M, Colombini B, Amedei A, Sofi F. Circadian rhythms, gut microbiota, and diet: Possible implications for health. *Nutr Metab Cardiovasc Dis*. 2023;33:1490-500. doi: 10.1016/j.numecd.2023.05.009. PubMed PMID: 37246076.
- Gutierrez Lopez DE, Lashinger LM, Weinstock GM, Bray MS. Circadian rhythms and the gut microbiome synchronize the host's metabolic response to diet. *Cell Metab*. 2021;33:873-87. doi: 10.1016/j.cmet.2021.03.015. PubMed PMID: 33789092.
- Neroni B, Evangelisti M, Radocchia G, Di Nardo G, Pantanella F, Villa MP, et al. Relationship between sleep disorders and gut dysbiosis: what affects what? *Sleep Med*. 2021;87:1-7. doi: 10.1016/j.sleep.2021.08.003. PubMed PMID: 34479058.
- Parkar SG, Kalsbeek A, Cheeseman JF. Potential Role for the Gut Microbiota in Modulating Host Circadian Rhythms and Metabolic Health. *Microorganisms*. 2019;7. doi: 10.3390/microorganisms7020041. PubMed PMID: 30709031; PubMed Central PMCID: PMC6406615.
- Hirotsu C, Tufik S, Andersen ML. Interactions between sleep, stress, and metabolism: From physiological to pathological conditions. *Sleep Sci*. 2015;8:143-52.

- doi: 10.1016/j.slscl.2015.09.002. PubMed PMID: 26779321; PubMed Central PMCID: PMC4688585.
- 19 Marwaha K, Cain R, Asmis K, Czaplinski K, Holland N, Mayer DCG, et al. Exploring the complex relationship between psychosocial stress and the gut microbiome: implications for inflammation and immune modulation. *J Appl Physiol* (1985). 2025;138:518-35. doi: 10.1152/japplphysiol.00652.2024. PubMed PMID: 39813028.
 - 20 Rusch JA, Layden BT, Dugas LR. Signalling cognition: the gut microbiota and hypothalamic-pituitary-adrenal axis. *Front Endocrinol (Lausanne)*. 2023;14:1130689. doi: 10.3389/fendo.2023.1130689. PubMed PMID: 37404311; PubMed Central PMCID: PMC10316519.
 - 21 Yang DF, Huang WC, Wu CW, Huang CY, Yang YSH, Tung YT. Acute sleep deprivation exacerbates systemic inflammation and psychiatry disorders through gut microbiota dysbiosis and disruption of circadian rhythms. *Microbiol Res*. 2023;268:127292. doi: 10.1016/j.micres.2022.127292. PubMed PMID: 36608535.
 - 22 Alruwaili NW, Alqahtani N, Alanazi MH, Alanazi BS, Aljrbua MS, Gatar OM. The effect of nutrition and physical activity on sleep quality among adults: a scoping review. *Sleep Science and Practice*. 2023;7:8. doi: 10.1186/s41606-023-00090-4.
 - 23 Theorell-Haglow J, Lemming EW, Michaelsson K, Elmstahl S, Lind L, Lindberg E. Sleep duration is associated with healthy diet scores and meal patterns: results from the population-based EpiHealth study. *J Clin Sleep Med*. 2020;16:9-18. doi: 10.5664/jcsm.8112. PubMed PMID: 31957658; PubMed Central PMCID: PMC7052994.
 - 24 Aziz T, Hussain N, Hameed Z, Lin L. Elucidating the role of diet in maintaining gut health to reduce the risk of obesity, cardiovascular and other age-related inflammatory diseases: recent challenges and future recommendations. *Gut Microbes*. 2024;16:2297864. doi: 10.1080/19490976.2023.2297864. PubMed PMID: 38174551; PubMed Central PMCID: PMC10773664.
 - 25 Kim B, Song A, Son A, Shin Y. Gut microbiota and epigenetic choreography: Implications for human health: A review. *Medicine (Baltimore)*. 2024;103:e39051. doi: 10.1097/MD.00000000000039051. PubMed PMID: 39029010; PubMed Central PMCID: PMC11398772.
 - 26 Reva K, Laranjinha J, Rocha BS. Epigenetic Modifications Induced by the Gut Microbiota May Result from What We Eat: Should We Talk about Precision Diet in Health and Disease? *Metabolites*. 2023;13. doi: 10.3390/metabo13030375. PubMed PMID: 36984815; PubMed Central PMCID: PMC10051796.
 - 27 Wu Q, Gao G, Kwok LY, Lv H, Sun Z. Insomnia: the gut microbiome connection, prospects for probiotic and postbiotic therapies, and future directions. *J Adv Res*. 2026;82:267-93. doi: 10.1016/j.jare.2025.07.005. PubMed PMID: 40651630; PubMed Central PMCID: PMC13000951.
 - 28 Morrison DJ, Preston T. Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism. *Gut Microbes*. 2016;7:189-200. doi: 10.1080/19490976.2015.1134082. PubMed PMID: 26963409; PubMed Central PMCID: PMC4939913.
 - 29 Archana, Gupta AK, Noumani A, Panday DK, Zaidi F, Sahu GK, et al. Gut microbiota derived short-chain fatty acids in physiology and pathology: An update. *Cell Biochem Funct*. 2024;42:e4108. doi: 10.1002/cbf.4108. PubMed PMID: 39228159.
 - 30 Lange O, Proczko-Stepaniak M, Mika A. Short-Chain Fatty Acids-A Product of the Microbiome and Its Participation in Two-Way Communication on the Microbiome-Host Mammal Line. *Curr Obes Rep*. 2023;12:108-26. doi: 10.1007/s13679-023-00503-6. PubMed PMID: 37208544; PubMed Central PMCID: PMC10250490.
 - 31 Magzal F, Even C, Haimov I, Agmon M, Asraf K, Shochat T, et al. Associations between fecal short-chain fatty acids and sleep continuity in older adults with insomnia symptoms. *Sci Rep*. 2021;11:4052. doi: 10.1038/s41598-021-83389-5. PubMed PMID: 33603001; PubMed Central PMCID: PMC7893161.
 - 32 O'Riordan KJ, Collins MK, Moloney GM, Knox EG, Aburto MR, Fulling C, et al. Short chain fatty acids: Microbial metabolites for gut-brain axis signalling. *Mol Cell Endocrinol*. 2022;546:111572. doi: 10.1016/j.mce.2022.111572. PubMed PMID: 35066114.
 - 33 Peng L, Li ZR, Green RS, Holzman IR, Lin J. Butyrate enhances the intestinal barrier by facilitating tight junction assembly via activation of AMP-activated protein kinase in Caco-2 cell monolayers. *J Nutr*. 2009;139:1619-25. doi: 10.3945/jn.109.104638. PubMed PMID: 19625695; PubMed Central PMCID: PMC2728689.
 - 34 Yan H, Ajuwon KM. Butyrate modifies

- intestinal barrier function in IPEC-J2 cells through a selective upregulation of tight junction proteins and activation of the Akt signaling pathway. *PLoS One*. 2017;12:e0179586. doi: 10.1371/journal.pone.0179586. PubMed PMID: 28654658; PubMed Central PMCID: PMC5487041.
- 35 Kim CH. Complex regulatory effects of gut microbial short-chain fatty acids on immune tolerance and autoimmunity. *Cell Mol Immunol*. 2023;20:341-50. doi: 10.1038/s41423-023-00987-1. PubMed PMID: 36854801; PubMed Central PMCID: PMC10066346.
 - 36 Dalile B, Van Oudenhove L, Vervliet B, Verbeke K. The role of short-chain fatty acids in microbiota-gut-brain communication. *Nat Rev Gastroenterol Hepatol*. 2019;16:461-78. doi: 10.1038/s41575-019-0157-3. PubMed PMID: 31123355.
 - 37 Church JS, Bannish JAM, Adrian LA, Rojas Martinez K, Henshaw A, Schwartz JJ. Serum short chain fatty acids mediate hippocampal BDNF and correlate with decreasing neuroinflammation following high pectin fiber diet in mice. *Front Neurosci*. 2023;17:1134080. doi: 10.3389/fnins.2023.1134080. PubMed PMID: 37123365; PubMed Central PMCID: PMC10130583.
 - 38 Banarase TA, Sammeta SS, Wankhede NL, Mangrulkar SV, Rahangdale SR, Aglawe MM, et al. Mitophagy regulation in aging and neurodegenerative disease. *Biophys Rev*. 2023;15:239-55. doi: 10.1007/s12551-023-01057-6. PubMed PMID: 37124925; PubMed Central PMCID: PMC10133433.
 - 39 Deng I, Corrigan F, Zhai G, Zhou XF, Bobrovskaya L. Lipopolysaccharide animal models of Parkinson's disease: Recent progress and relevance to clinical disease. *Brain Behav Immun Health*. 2020;4:100060. doi: 10.1016/j.bbih.2020.100060. PubMed PMID: 34589845; PubMed Central PMCID: PMC8474547.
 - 40 Hou K, Wu ZX, Chen XY, Wang JQ, Zhang D, Xiao C, et al. Microbiota in health and diseases. *Signal Transduct Target Ther*. 2022;7:135. doi: 10.1038/s41392-022-00974-4. PubMed PMID: 35461318; PubMed Central PMCID: PMC9034083.
 - 41 Solanki R, Karande A, Ranganathan P. Emerging role of gut microbiota dysbiosis in neuroinflammation and neurodegeneration. *Front Neurol*. 2023;14:1149618. doi: 10.3389/fneur.2023.1149618. PubMed PMID: 37255721; PubMed Central PMCID: PMC10225576.
 - 42 Wang Z, Chen WH, Li SX, He ZM, Zhu WL, Ji YB, et al. Gut microbiota modulates the inflammatory response and cognitive impairment induced by sleep deprivation. *Mol Psychiatry*. 2021;26:6277-92. doi: 10.1038/s41380-021-01113-1. PubMed PMID: 33963281.
 - 43 Zhang M, Zhang M, Kou G, Li Y. The relationship between gut microbiota and inflammatory response, learning and memory in mice by sleep deprivation. *Front Cell Infect Microbiol*. 2023;13:1159771. doi: 10.3389/fcimb.2023.1159771. PubMed PMID: 37293204; PubMed Central PMCID: PMC10244646.
 - 44 Zhao M, Chu J, Feng S, Guo C, Xue B, He K, et al. Immunological mechanisms of inflammatory diseases caused by gut microbiota dysbiosis: A review. *Biomed Pharmacother*. 2023;164:114985. doi: 10.1016/j.biopha.2023.114985. PubMed PMID: 37311282.
 - 45 Fang H, Yao T, Li W, Pan N, Xu H, Zhao Q, et al. Efficacy and safety of fecal microbiota transplantation for chronic insomnia in adults: a real world study. *Front Microbiol*. 2023;14:1299816. doi: 10.3389/fmicb.2023.1299816. PubMed PMID: 38088972; PubMed Central PMCID: PMC10712199.
 - 46 Kim CS, Cha L, Sim M, Jung S, Chun WY, Baik HW, et al. Probiotic Supplementation Improves Cognitive Function and Mood with Changes in Gut Microbiota in Community-Dwelling Older Adults: A Randomized, Double-Blind, Placebo-Controlled, Multicenter Trial. *J Gerontol A Biol Sci Med Sci*. 2021;76:32-40. doi: 10.1093/gerona/glaa090. PubMed PMID: 32300799; PubMed Central PMCID: PMC7861012.
 - 47 Salonen T, Jokinen E, Satokari R, Lahtinen P. Randomized, double-blinded, placebo-controlled pilot study: efficacy of faecal microbiota transplantation on chronic fatigue syndrome. *J Transl Med*. 2023;21:513. doi: 10.1186/s12967-023-04227-y. PubMed PMID: 37516837; PubMed Central PMCID: PMC10386223.
 - 48 Sanborn V, Aljumaah M, Azcarate-Peril MA, Gunstad J. Examining the cognitive benefits of probiotic supplementation in physically active older adults: A randomized clinical trial. *Appl Physiol Nutr Metab*. 2022;47:871-82. doi: 10.1139/apnm-2021-0557. PubMed PMID: 35617704.
 - 49 Skjevling L, Goll R, Hanssen HM, Johnsen PH. Faecal microbiota transplantation (FMT) in Norwegian outpatients with mild to severe myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS): protocol for a 12-month

- randomised double-blind placebo-controlled trial. *BMJ Open*. 2024;14:e073275. doi: 10.1136/bmjopen-2023-073275. PubMed PMID: 38858151; PubMed Central PMCID: PMC11168185.
- 50 Tanaka A, Sanada K, Miyaho K, Tachibana T, Kurokawa S, Ishii C, et al. The relationship between sleep, gut microbiota, and metabolome in patients with depression and anxiety: A secondary analysis of the observational study. *PLoS One*. 2023;18:e0296047. doi: 10.1371/journal.pone.0296047. PubMed PMID: 38117827; PubMed Central PMCID: PMC10732403.
 - 51 Grosicki GJ, Riemann BL, Flatt AA, Valentino T, Lustgarten MS. Self-reported sleep quality is associated with gut microbiome composition in young, healthy individuals: a pilot study. *Sleep Med*. 2020;73:76-81. doi: 10.1016/j.sleep.2020.04.013. PubMed PMID: 32795890; PubMed Central PMCID: PMC7487045.
 - 52 Haimov I, Magzal F, Tamir S, Lalzar M, Asraf K, Milman U, et al. Variation in Gut Microbiota Composition is Associated with Sleep Quality and Cognitive Performance in Older Adults with Insomnia. *Nat Sci Sleep*. 2022;14:1753-67. doi: 10.2147/NSS.S377114. PubMed PMID: 36225322; PubMed Central PMCID: PMC9550024.
 - 53 Zhang Q, Yun Y, An H, Zhao W, Ma T, Wang Z, et al. Gut Microbiome Composition Associated With Major Depressive Disorder and Sleep Quality. *Front Psychiatry*. 2021;12:645045. doi: 10.3389/fpsyt.2021.645045. PubMed PMID: 34093266; PubMed Central PMCID: PMC8175648.
 - 54 Ahmadi S, Taghizadieh M, Mehdizadehfard E, Hasani A, Khalili Fard J, Feizi H, et al. Gut microbiota in neurological diseases: Melatonin plays an important regulatory role. *Biomed Pharmacother*. 2024;174:116487. doi: 10.1016/j.biopha.2024.116487. PubMed PMID: 38518598.
 - 55 Gao K, Mu CL, Farzi A, Zhu WY. Tryptophan Metabolism: A Link Between the Gut Microbiota and Brain. *Adv Nutr*. 2020;11:709-23. doi: 10.1093/advances/nmz127. PubMed PMID: 31825083; PubMed Central PMCID: PMC7231603.
 - 56 Jenkins TA, Nguyen JC, Polglaze KE, Bertrand PP. Influence of Tryptophan and Serotonin on Mood and Cognition with a Possible Role of the Gut-Brain Axis. *Nutrients*. 2016;8. doi: 10.3390/nu8010056. PubMed PMID: 26805875; PubMed Central PMCID: PMC4728667.
 - 57 Mhanna A, Martini N, Hmaydoosh G, Hamwi G, Jarjanazi M, Zaifiah G, et al. The correlation between gut microbiota and both neurotransmitters and mental disorders: A narrative review. *Medicine (Baltimore)*. 2024;103:e37114. doi: 10.1097/MD.00000000000037114. PubMed PMID: 38306525; PubMed Central PMCID: PMC10843545.
 - 58 Suda K, Matsuda K. How Microbes Affect Depression: Underlying Mechanisms via the Gut-Brain Axis and the Modulating Role of Probiotics. *Int J Mol Sci*. 2022;23. doi: 10.3390/ijms23031172. PubMed PMID: 35163104; PubMed Central PMCID: PMC8835211.
 - 59 Chen Y, Xu J, Chen Y. Regulation of Neurotransmitters by the Gut Microbiota and Effects on Cognition in Neurological Disorders. *Nutrients*. 2021;13. doi: 10.3390/nu13062099. PubMed PMID: 34205336; PubMed Central PMCID: PMC8234057.
 - 60 Hamamah S, Aghazarian A, Nazaryan A, Hajnal A, Covasa M. Role of Microbiota-Gut-Brain Axis in Regulating Dopaminergic Signaling. *Biomedicines*. 2022;10. doi: 10.3390/biomedicines10020436. PubMed PMID: 35203645; PubMed Central PMCID: PMC8962300.
 - 61 Pires L, Gonzalez-Paramás AM, Heleno SA, Calhella RC. Gut Microbiota as an Endocrine Organ: Unveiling Its Role in Human Physiology and Health. *Applied Sciences*. 2024;14:9383. doi:10.3390/app14209383.
 - 62 Siegel JM. The neurotransmitters of sleep. *J Clin Psychiatry*. 2004;65:4-7. PubMed PMID: 15575797; PubMed Central PMCID: PMC8761080.
 - 63 Wang Y, Qiao H, Zhang Y. Changes of Dopamine and Tyrosine Hydroxylase Levels in the Brain of Germ-free Mice. *Iran J Biotechnol*. 2023;21:e2798. doi: 10.30498/ijb.2022.236732.2798. PubMed PMID: 36811101; PubMed Central PMCID: PMC9938933.
 - 64 Yassin LK, Nakhal MM, Alderei A, Almehairbi A, Mydeen AB, Akour A, et al. Exploring the microbiota-gut-brain axis: impact on brain structure and function. *Front Neuroanat*. 2025;19:1504065. doi: 10.3389/fnana.2025.1504065. PubMed PMID: 40012737; PubMed Central PMCID: PMC11860919.
 - 65 Bjorvatn B, Gronli J, Hamre F, Sorensen E, Fiske E, Bjorkum AA, et al. Effects of sleep deprivation on extracellular serotonin in hippocampus and frontal cortex of the rat. *Neuroscience*. 2002;113:323-30. doi: 10.1016/s0306-4522(02)00181-1. PubMed PMID: 12000000.

- 12127089.
- 66 Gruenbaum BF, Merchant KS, Zlotnik A, Boyko M. Gut Microbiome Modulation of Glutamate Dynamics: Implications for Brain Health and Neurotoxicity. *Nutrients*. 2024;16. doi: 10.3390/nu16244405. PubMed PMID: 39771027; PubMed Central PMCID: PMC11677762.
- 67 Guo J, Guo J, Rao X, Zhang R, Li Q, Zhang K, et al. Exploring the pathogenesis of insomnia and acupuncture intervention strategies based on the microbiota-gut-brain axis. *Front Microbiol*. 2024;15:1456848. doi: 10.3389/fmicb.2024.1456848. PubMed PMID: 39364160; PubMed Central PMCID: PMC11446747.
- 68 Varintra P, Anwar S, Shih SC, Liu IY. The role of the GABAergic system on insomnia. *Tzu Chi Med J*. 2024;36:103-9. doi: 10.4103/tcmj.tcmj_243_23. PubMed PMID: 38645778; PubMed Central PMCID: PMC11025592.
- 69 Zheng P, Zeng B, Liu M, Chen J, Pan J, Han Y, et al. The gut microbiome from patients with schizophrenia modulates the glutamate-glutamine-GABA cycle and schizophrenia-relevant behaviors in mice. *Sci Adv*. 2019;5:eaau8317. doi: 10.1126/sciadv.aau8317. PubMed PMID: 30775438; PubMed Central PMCID: PMC6365110.
- 70 Dahlin M, Elfving A, Ungerstedt U, Amark P. The ketogenic diet influences the levels of excitatory and inhibitory amino acids in the CSF in children with refractory epilepsy. *Epilepsy Res*. 2005;64:115-25. doi: 10.1016/j.epilepsyres.2005.03.008. PubMed PMID: 15961283.
- 71 Wu M, Zhang X, Feng S, Freda SN, Kumari P, Dumrongprechachan V, et al. Dopamine pathways mediating affective state transitions after sleep loss. *Neuron*. 2024;112:141-54. doi: 10.1016/j.neuron.2023.10.002. PubMed PMID: 37922904; PubMed Central PMCID: PMC10841919.